

Vespasiano, G.^{1,2}, Cianflone, G.^{1,2,}, Marini L.³, De Rosa, R.¹, Polemio, M.⁴, Walraevens, K.⁵, Vaselli, O.^{6,7}, Pizzino, L.⁸, Cinti, D.⁸, Capecchiacci, F.^{6,7,9}, Barca D.¹, Dominici, R.^{1,2} and Apollaro, C.¹*

* Corresponding author: giuseppe.cianflone@unical.it

Abstract

This work pursues the hydro-geochemical and isotopic characterization of the complex groundwater system of the Gioia Tauro Plain, one of the most important industrialized and agricultural coastal areas of southern Italy. The anthropic pressure exposes the water resources at risk of depletion and quality degradation making the plain groundwater a system of high scientific and social interest.

The plain is characterised by a shallow aquifer, mostly recharged by local rains and a deep aquifer apparently less influenced by local precipitation. Both aquifers are mainly Ca-HCO₃ waters except for localized sectors where Na-HCO₃, Na-Cl and Ca-SO₄ waters are present. In deep aquifer, both prolonged interaction with sedimentary rocks, mainly deriving from the erosion of crystalline rocks, and direct cation exchange represent the primary factors controlling the formation of Na-HCO₃ waters. Mixing processes between these waters and either connate brine and/or deep thermal waters contribute to the formation of isolated high salinity Na-Cl-rich waters. In shallow aquifer, inputs of N-rich sewage and agriculture-related contaminants, and SO_x emissions in proximity of the harbour are responsible of the increasing nitrate and sulphate concentrations, respectively. The Cl/Br and NO₃/Cl ratios highlight contamination mainly linked to agricultural activities and contribution of wastewater.

Along the northern boundary, the warmest groundwater (Na-Cl[SO₄]) were found close to a bend of the main strike-slip fault system, locally favouring the rising of B- and Li-rich deep waters, testifying the influence of geological-structural features on deep water circulation.

Despite the high-water demand, a direct marine intrusion is localized in a very restricted area, where we observed an incipient groundwater-seawater mixing (seawater contribution $\leq 7\%$). The qualitative and quantitative conditions of the shallow aquifer still have acceptable levels because of the relatively high recharge inflow. A reliable hydrogeochemical conceptual model, able to explain the compositional variability of the studied waters, is proposed.

Keywords: Coastal aquifers - hydrogeochemistry - water isotopes – conceptual hydrogeochemical model – Calabria – Southern Italy

1. Introduction

During the last years, the sustainable management of coastal plains have received significant attention worldwide (e.g., Boughriba & Jilali, 2018; Matiatos et al., 2018; Vespasiano et al., 2015a, 2019; Mastrocicco & Colombani, 2020; Polemio & Zuffianò, 2020; Cianflone et al., 2021; Sarker et al., 2021; Nisi et al., 2022). Coastal plains show an important economic, social, and biological value as they represent the link between marine and terrestrial environments (Martínez et al., 2007). In response to this, coastal plains have always been considered strategic sectors for human settlement hosting most of the world's megacities with the relative issues due to the intensive use of the territory (McGranahan et al., 2007). In these areas, the intensive exploitation of groundwater reduces freshwater outflow to the sea, forming a local water table depression and inland seawater migration (Masciopinto et al., 2006; Mjemah et al., 2009; Van Camp et al., 2014). This phenomenon, named seawater intrusion, is a global concern and represents one of the most widespread processes that degrade groundwater quality and limit its utilization (Nisi et al., 2013; Alfarrach and Walraevens, 2018; Vespasiano et al., 2019; Boumaiza et al., 2020; Han et al., 2021; Kammoun et al., 2021). Coupled with a continuing sea level rise due to global warming, coastal aquifers are expected to be threatened in a relatively near future (Alfarrach and Walraevens, 2018). Consequently, there has been an increasing interest in evaluating the extent of seawater intrusion in response to overexploitation and sea level rise (Barlow and Reichard, 2010; Custodio, 2010; Polemio and Walraevens, 2019). Seawater intrusion phenomena have been reported in many coastal areas of the United States of America (Barlow, 2003; Langman and Ellis, 2010), Africa (El Halimi et al., 2001; Steyl and Dennis, 2010; Van Camp et al., 2013; Walraevens et

al., 2015; Alfarrach and Walraevens, 2018) and, to a lesser extent, of Mexico (Cardoso, 1993), South America (Bocanegra et al., 2010), Saudi Arabia (Alshehri et al., 2021) and Australia (Werner, 2010). Also in Italy, studies focused on the sustainable management and characterization of the main processes linked to marine intrusion have undergone a considerable increase (e.g., Capaccioni et al., 2005). From the northern-central to the southernmost coasts of the Adriatic and Tyrrhenian Seas, as well as in Sardinia and Sicily, several areas were recognized to be at risk of salinization (e.g., Barrocu, 2003; Grassi et al. 2007; Brozzo et al., 2011; Mollema et al., 2013; Felisa et al., 2013; Iyalomhe et al., 2015; Franceschini and Signorini, 2016; Vespasiano et al., 2016; Vespasiano et al., 2019; Muzzillo et al., 2020; Bellafiore et al., 2021; Giambastiani et al., 2021; Mastrocicco et al., 2021; Cianflone et al., 2021, 2022), similarly to what is occurring in other coastal areas of the Mediterranean Sea (e.g., Alcalá and Custodio, 2008; De Montety et al., 2008; Custodio, 2010; Mastrocicco and Colombani, 2020).

Complex geological setting, mixing between marine and connate waters, ion exchange processes and natural and/or human-induced inputs (due to industrial and agricultural activities) represent other conditions that can modify groundwater composition of coastal plains (Mollema et al., 2013; Mtoni et al., 2013; Carreira et al., 2014; Corniello et al. 2015; Abu Al Nacem et al., 2019; Vespasiano et al., 2019; Mastrocicco et al., 2021). In these contexts, sources and driving mechanisms are to be identified and characterized by a multidisciplinary approach (including geological, hydrogeological, geochemical, and geophysical investigations), to produce reliable conceptual and numerical models whose results can be used for steering suitable preventive actions. Water geochemistry and isotopes (e.g., $^{18}\text{O}/^{16}\text{O}$ and $^2\text{H}/^1\text{H}$ ratios) are useful to: i) identify the main ongoing water-rock interaction processes, leading to the formation of the different geochemical facies (Vespasiano et al., 2019); ii) evaluate freshwater-saltwater interactions as well as consequent ion exchange and oxidation–reduction reactions (e.g., Zarei et al., 2012; Sudaryanto and Naily, 2018); iii) contribute to the management of the groundwater resource (e.g., Menichini and Doveri, 2020; Carrión-Mero et al., 2021); iv) model transport pathways and fate of saline water (e.g., Ganyaglo et al., 2017; Wua et al., 2020); v) evaluate the environmental impact induced by human activities, if any, and vi) assess groundwater quality (e.g., Apollaro et al., 2019; Vardè et al., 2019; Apollaro et al., 2020; Figoli et al., 2020; Alshehri et al., 2021; Mastrocicco et al., 2021; Apollaro et al., 2022). In coastal plains, a detailed geochemical characterization of the aforementioned processes represents the primary objective to promote sustainable management and intervene purposefully for the resolution and/or mitigation of possible problems (Fehdi et al., 2009; Bompoti et al., 2015; Mallick et al., 2018).

This work is aimed at investigating the Gioia Tauro Plain (GTP) groundwater system. GTP is located along the southwestern coast of the Calabria Region and is one of the most important industrialized and coastal agricultural areas of southern Italy, with an extension of about 500 km². The plain is inhabited by about 160,000 people distributed in towns, villages, and rural areas. It is characterized by a highly developed agricultural sector and two thermoelectric industries (waste-to-energy and thermoelectric power plants). Furthermore, GTP hosts two strategic infrastructures such as the Gioia Tauro Harbor (the largest transshipment terminal of Italy and one of the most important container traffic hubs in the Mediterranean basin) and the A2 “Mediterranea” highway (the main road line of the Region).

Despite the crucial role and the presence of relevant infrastructures, GTP still presents portions completely devoid of basic services, such as sewers or connections to aqueducts for domestic use. The lack of such services and the well-developed anthropic activities expose the GTP groundwater resources at risk of depletion and quality degradation. Furthermore, GTP shows a very articulated geological and structural framework that locally favour the development of waters with chemical characteristics not fully explainable by typical water-rock interaction processes occurring in the sedimentary aquifers and/or seawater intrusion (Pizzino et al., 2004; Cianflone et al., 2021). In addition to the ongoing

anthropic pressure (mainly due to intensive farming), this issue makes the GTP groundwater a system of high scientific and social interest.

This work has as main target that to improve the hydrogeological and geochemical knowledge of the GTP groundwater systems by providing new geochemical and isotopic data to i) assess the main processes affecting the main GTP aquifers and ii) establish the relationship between natural and anthropogenic sources that concur to the observed water chemistry variability and iii) propose a conceptual model of fluid circulation.


2. Geological, structural, and hydrogeological settings

GTP is a NNE-SSW oriented half-graben (Figure 1a) of Upper Pliocene-Pleistocene age (Monaco and Tortorici, 2000), bounded to the east by the NNE-SSW Cittanova Fault (CF_NNE-SSW_W) [*fault acronym_strike_dip*] (Figure 1a and Figure 1Sa), a high-angle, west-dipping, master normal fault and juxtaposing the crystalline bedrock and the GTP sedimentary infill (Jacques et al., 2001; Tripodi et al., 2018). CF was responsible for both the 1783 catastrophic earthquake (Mw = 7.0) and the seismic event that occurred in the 4th century AD (Galli and Peronace, 2015; Galli and Bosi, 2002). Other three faults parallel to CF occur in the study area: the west dipping Sant'Eufemia-Laureana Fault (SLF_NNE-SSW_W) and the Palmi-Gioia Tauro Fault (PGF_NNE-SSW_W) and an east dipping tectonic structure occupying an intermediate position between SLF and PGF: the Rosarno-Palmi Fault (RPF_NNE-SSW_E) (Tripodi et al., 2018). Furthermore, two other faults, orthogonal to CF, occur: (i) to the North, the Nicotera-Gioiosa Fault Zone (NGFZ_NW-SE_S), NW-SE striking and with right strike-slip kinematics, (ii) to the South, the Palmi-Locri Fault Zone (PLFZ_NW-SE_N), of NW-SE direction and a main left strike-slip kinematics (Tripodi et al., 2018 and references therein). According to Loreto et al. (2019), the GTP depression represents the emerged portion of the Gioia Tauro Basin, extending offshore in the Tyrrhenian Sea, while NGFZ is the subaerial part of an important tectonic structure continuing for some tens of km in the Tyrrhenian Sea and delimiting to the south the Capo Vaticano Ridge.

The eastern and northern sides of the plain are bounded by the Serre Massif, a complete continental crustal section made up of Variscan metamorphic and plutonic rocks (Cirrincione et al., 2015). The Serre Batholith crops out along the eastern boundary and includes two-mica porphyritic granodiorites and granites, two-mica granodiorites and granites, and biotite granodiorites (Figure 1a). The Capo Vaticano Promontory is a sector of the Serre Massif separated from the main body by the Mesima graben. It bounds the NW-sector of the plain and is made up of qz-diorites and tonalites, and a migmatitic border zone. To the south of Cittanova town, PLFZ_NW-SE_N separates the Serre Massif from the Aspromonte Massif (Cirrincione et al., 2015). In this sector, the latter consists of migmatitic paragneisses, which outcrop in the catchment of the Petrace River. The GTP southwestern side is bounded by the Palmi High, characterized by crystalline rocks (migmatitic paragneiss, foliated tonalite and augen gneiss) covered by Tortonian-Messinian siliciclastic and carbonate deposits (Cirrincione et al., 2015).

The GTP basin is filled by a sedimentary succession starting with upper Miocene siliciclastic and carbonate deposits, which unconformably cover the crystalline bedrock (Figure 1a). The succession passes upward to thick (~600 m) Plio-Pleistocene marine sediments (Jacques et al., 2001), which include, at their bottom, Pliocene clayey and silty deposits, hundred meters thick (Jacques et al., 2001), called "Trubi Formation" (ISPRA, 2016), and characterized in the upper part by the presence of a pumice-rich horizon (De Rosa et al., 2008). The Trubi Formation is covered, through a slight angular unconformity, by the Late Pliocene-Pleistocene Calcareni di Vinco, consisting of cross-bedded sands and calcarenites (Jacques et al., 2001; ISPRA, 2016), which are considered as the infilling of the Siderno paleo-strait (Longhitano et al., 2012). The Late Pleistocene gravel and sand alluvial fan deposits of the Taurianova Synthem cover the Calcareni di Vinco through an erosive contact, in the eastern side of the plain (ISPRA, 2016). The sedimentary infill of the GTP is

150 closed by Late Pleistocene–Holocene sand- to gravel-terraced marine deposits (up to 20 m thick) and Holocene terrains
 151 where ancient aeolian inland dunes, recent alluvial deposits (in the eastern and middle sectors), and coastal deposits (in
 152 the western sector) were recognized.
 153



154
 155 **Figure 1** – (a) Simplified geological map of the Gioia Tauro area (modified after [Cianflone et al., 2021](#)) showing location and ID
 156 ([Stuyfzand, 1986](#)) of water samples (red symbols: deep aquifer; black symbols: shallow aquifer); (b) schematic stratigraphic column
 157 (not to scale) showing the deep (HU1) and shallow (HU3) aquifers and the aquitard (HU2) recognized in the study area. NGFZ:
 158 Nicotera-Gioiosa Fault Zone; PGF: Palmi-Gioia Tauro Fault; RPF: Rosarno-Palmi Fault; SLF: Sant'Eufemia-Laureana Fault; CF:
 159 Cittanova Fault; PLFZ: Palmi-Locri Fault Zone.
 160

161 From a hydrogeological point of view, the GTP sub-surface hosts (i) a deep aquifer hosted in the Late Miocene succession
162 (HU1) and characterised by artesian condition, (ii) an aquitard (HU2) made up of Pliocene clayey and silty deposits and
163 (iii) a shallow phreatic aquifer corresponding to the Late Pleistocene and Holocene marine and alluvial sediments, with
164 hydraulic conductivity ranging from 10^{-4} to 10^{-5} m/s, somewhere with peaks of 10^{-3} m/s (HU3) (Figure 1b). A peculiar
165 hydrogeological setting characterizes the Scinà area (Figure 1a) where the shallow aquifer (HU3), consisting of fluvial
166 and deltaic sediments of the Petrace River and coastal deposits, lies directly over the metamorphic basement (Cianflone
167 et al., 2021). The occurrence of deeper water circuits within the crystalline basement is indicated by both the Galatro
168 thermal springs (Apollaro et al., 2019; Randazzo et al., 2022), in the north-eastern border of the study area (Figure 1a),
169 and B- and Li-rich waters along the NGFZ_ NW-SE_ S (Pizzino et al., 2004). The groundwater flow in the shallow aquifer
170 (Figure 2Sa) has a prevalent SE-NW direction.

171 The GTP comprises four main catchments (Figures 2Sb): (i) the Mesima basin (extension of about 815 km²), in the
172 northern sector with a SW, W and, subordinately, NW drainage. The basin drains the Serre Massif and, partially, the
173 Aspromonte Massif with the tributary Metramo River; (ii) the Budello basin, in the central portion, showing a W-NW
174 drainage, a small catchment and a torrential watercourse; (iii) the Petrace basin, located in the southern sector, showing a
175 N and NW drainage direction from the Aspromonte Massif (extension of about 420 km²) and iv) the small (about 20 km²)
176 San Giovanni basin, in the northernmost side of the plain with a main NS flow direction draining the Capo Vaticano
177 promontory.

178 The mean annual rainfall distribution was analysed according to Cianflone et al. (2015a). Data from 26 rain gauges
179 (Regione Calabria – ARPACAL - <https://www.cfd.calabria.it>) were used to define the correlation between the Average
180 Annual Rainfall (AAR) and Altitude (ALT) ($AAR = 1.0409 \times ALT + 846.08$, $R^2 = 0.7$). Then, we converted a DTM 20×20 m
181 (Geo-portale Nazionale - <http://www.pcn.minambiente.it/mattm>) from raster to a point shapefile (one point for each cell)
182 with associated altitude data. For each point, the previously obtained correlation was applied. Finally, the punctual data
183 were processed using the ordinary kriging as interpolation technique to obtain a continuous spatial reconstruction of the
184 average annual rainfall (Figure 2Sb), which is characterized by a mean annual value of 1274 mm.

185

186 3. Methods

187 3.1. Sampling and analytical methods

188 The hydrogeochemical survey at GTP was carried out from March to October 2021. A total of 114 water points, including
189 wells, rivers, and springs were selected for water level measurement, physicochemical field measurements and
190 geochemical (major and trace solutes) and isotopic ($\delta^{18}\text{O}$ and $\delta^2\text{H}$) laboratory analyses (Table 1S). The water samples
191 were identified with a progressive number preceded by a suffix: PG: well; FG: river and SG: spring (Figure 1Sa). The
192 well groundwater samples were collected during pumping and the water level measurements were performed beforehand
193 pumping (for the elaboration of groundwater levels see Cianflone et al., 2021). Wells were previously purged with
194 international standard criteria (generally low flow and pumping of 3-4 volumes of well water column); and the samples
195 were collected once the physicochemical parameters stabilized. The 114 samples included: 78 samples of the shallow
196 aquifer (using wells from 5 to 100 m depth), 21 samples from the deep confined aquifer (using artesian wells from 90 to
197 436 m depth), 9 samples from rivers (FG1, FG2, and FG3 of Budello River; FG4, FG5, and FG6 of Mesima River; FG7
198 of Metramo River; FG8 and FG9 Petrace River). Furthermore, 6 samples (5 from wells drilled in the metamorphic-
199 crystalline basement and 1 from a spring), representative of the main recharge areas, were collected: Palmi High (PG94),
200 Aspromonte massif (PG36) and Capo Vaticano High (PG16, PG17, SG4 and PG71) (Figure 1Sa). The three recharge

201 areas were treated separately, given the compositional differences of the outcropping lithotypes that could affect the
 202 composition of the recharge waters.

203 The assignment of each well to the shallow (HU3) or deep aquifer (HU1) was carried out by comparing the well depth
 204 with the reconstructed stratigraphy, considering artesian or phreatic evidence.

205 It was generally assumed that groundwater samples are mainly representative of the deepest part of wells, if they were
 206 bored to tap groundwater and not for monitoring purposes (boring local companies usually are paid drill up to find enough
 207 groundwater).

208 Intrinsically unstable parameters (total alkalinity, temperature, pH, and Eh) and electrical conductivity (EC) were
 209 measured in the field by means of portable instruments (Hanna instruments - *HI9829*). Two pH buffers, with nominal pH
 210 values of 4.01 and 7.01 at 25 °C, were used for pH calibration at each sampling site. The ZoBell's solution ([Nordstrom, 1997](#))
 211 was utilized to calibrate the mV-meter for Eh measurement. Total alkalinity was determined by acidimetric titration,
 212 using HCl 0.05N as titrating agent and methyl-orange as indicator ([Nisi et al., 2019](#); [Vaselli et al., 2021](#)). All water
 213 samples were filtered in situ through a membrane with a 0.45 µm pore (e.g., [Kennedy and Zellweger, 1974](#); [Laxen and Chandler, 1982](#)).
 214 Water samples for the determination of anions were stored without additional treatments, whereas
 215 samples for the determination of cations, trace (B, Li, Al, Fe and Mn) elements, and SiO₂ were acidified by adding
 216 suprapur acid (1% HNO₃). All the samples were stored in polyethylene bottles, previously washed with dilute HNO₃ and
 217 rinsed with Milli-Q demineralized water. Blank solutions were prepared in the field using demineralized water and
 218 following the same procedure. All the water samples were kept at 4 °C and dark conditions before analysis. To measure
 219 the δ²H and δ¹⁸O values, 50 ml of water was transferred into tightly sealed screw-cap bottles without filtering. Waterproof
 220 tape was bound around the cap to ensure that no evaporation took place.

221 The main solutes were determined by High Performance Liquid Chromatography (HPLC) by using a Thermo Scientific
 222 Dionex™ ICS-1100 equipped with Dionex IOnPac AS23 and Dionex IonPac CS12A columns for the determination of
 223 anionic (F⁻, Cl⁻, Br⁻, SO₄²⁻ and NO₃⁻) and cation (K⁺, Na⁺, Li⁺, Ca²⁺, NH₄⁺ and Mg²⁺) species, respectively, whereas, trace
 224 elements (B, Al, Fe and Mn) were determined with a quadrupole inductively coupled plasma-mass spectrometer which
 225 utilizes a dynamic reaction cell for interference removal (ICP-MS, PerkinElmer/SCIEX, ELAN DRC-e). On the same
 226 day of collection, dissolved reactive SiO₂ was measured by VIS spectrophotometry upon reaction with ammonium
 227 molybdate in acid media (and treatment with oxalic acid) to form a yellow silicomolybdate complex, whose absorbance
 228 was read at 410 nm (see [Nollet and De Gelder, 2007](#) for further details). Data quality for major components was evaluated
 229 by calculating the electroneutrality parameter (EP; [Appelo and Postma, 2005](#)) charge balance, as follows:

$$EP = [(\Sigma cat - \Sigma an) / (\Sigma cat + \Sigma an)] * 100$$

233 The sum of cations (Σcat) and anions (Σan) are in meq/L. The EP was always less than ±5 %. Precision and accuracy for
 234 minor and trace elements was checked against the NIST1643f standard reference solution. Deviations from the certified
 235 concentrations resulted to be lower than 5%. The δ²H and δ¹⁸O (both in ‰ V-SMOW) values were measured in the
 236 Laboratories of the Stable Isotopes at the University of Florence by a Picarro L2130-i isotope analyzer, which utilise a
 237 cavity ring-down spectroscopy (CRDS) technology. The internal standards are originally inter-calibrated by using the
 238 IAEA international standards. During the instrumental setup, the possible “salinity” effects were verified. In particular,
 239 at MilliQ water increasing amounts of suprapur (dried) NaCl were added up to salinity similar or even higher than that of
 240 seawater were achieved. During these tests no significant differences were recorded. The vaporization system indeed uses
 241 a protective septum to reduce the deposition of salt in the vaporization chamber. Additionally, when dealing with saline

samples we test the accuracy of our instrument with an internal seawater from the Mediterranean Sea that has been cross-analyzed with other isotopic labs, e.g. the INGV Vesuvian Observatory. It is also to mention that Picarro L2130-i has 7 positions for each line, in each line one position is dedicated to MilliQ water to control any drifts. Before each run, the internal standards are analyzed 3 to 4 times and then, every 12 samples and at the end of the any analytical session to verify and optimize the calibration. Each sample is analyzed seven times and the first 3 measurements are excluded from the post-process program to avoid any memory effects. Measurement precision, based on the repeated analysis of in-house standards, was found to be $\pm 0.67\%$ for $\delta^2\text{H}$ and $\pm 0.09\%$ for $\delta^{18}\text{O}$. In terms of accuracy, the error is $\pm 0.34\%$ for $\delta^2\text{H}$ and $\pm 0.055\%$ for $\delta^{18}\text{O}$.

3.2. Data processing

The final aim of the data interpretation is to elaborate a site-specific conceptual model of GTP based on geochemical/isotopic data as well as on geological and hydrogeological evidence. Thus, the study area was divided in 6 main sectors (**Figure 1Sb**), as a function of the lithological-structural framework: Sector A (eastern portion-recharge area), B (northern portion across the NGFZ_ NW-SE_S), C (north-central portion), D (south-central portion), E (Scinà Area) and F (southern portion across the PLFZ_ NW-SE_N).

The geochemical data collected during this work were also compared with those of the Galatro thermal waters ([Pizzino et al., 2004](#); [Apollaro et al., 2019](#)) since the latter likely represent the discharge at surface of the main deep thermal system present in the study area.

The geochemical data were interpreted according to consolidated procedures, such as: (i) the Durov diagram ([Durov 1948](#) – **Figure 2**); (ii) the Total Ionic Salinity (TIS) diagram of $(\text{Ca}^{2+}+\text{Mg}^{2+})$ vs. $(\text{Na}^{+}+\text{K}^{+})$ (**Figure 3a** and **3b**), in which the iso-TIS lines are drawn for reference ([Apollaro, 2019](#); [Vespasiano et al., 2021a](#); [Vespasiano et al., 2021b](#)); (iii) binary plots between mobile and conservative components. Saturation indexes (SI) were calculated for selected mineral phases using the PHREEQC interactive software, *version 3.1.1* ([Parkhurst and Appelo, 2013](#)) and the LLNL thermodynamic database (**Table 1S**).

In the Durov diagram, the result is a representation of nine fields (from ^(a) to ⁽ⁱ⁾ in **figure 2**) that allows to separate properly the four main chemical facies generally recognized in coastal areas [$\text{Na-HCO}_3^{(c)}$, $\text{Na-Cl}^{(i)}$, $\text{Ca-HCO}_3^{(a)}$ and $\text{Ca-Cl}^{(g)}$] as well as the mixed chemical types occupying intermediate positions ([Petalas et al., 2009](#)).

A complete description of the GTP waters was performed according to their chemical and physical features by adopting a simplified version of the classification approach proposed by [Stuyfzand \(1989\)](#). Water classification by [Stuyfzand \(1989\)](#) is commonly applied to ground and surface waters occurring in coastal areas and used to identify cation exchange reactions and specific geochemical processes, which possibly control their water chemistry (e.g., [Giménez-Forcada et al., 2010](#); [Marconi et al., 2011](#); [Vandenbohede and Lebbe, 2012](#); [Mendizabal et al., 2012](#); [Mollema et al., 2013](#); [Sarker et al., 2021](#); [Alfarrah and Walraevens, 2018](#)).

The approach of [Stuyfzand \(1989\)](#) implies the step-by-step determination of the main type, type, subtype, and class (Base Exchange Index - BEX) of each water sample. Each of these four levels of classification is recorded in the total code (and name) of the water type. The main type of water is based on Cl^- concentration (in mg L^{-1}). These are G = very oligohaline: <5 ; g = oligohaline: 5-30; F = fresh: 30-150; f = fresh-brackish: 150-300; B = brackish: 300-1000; b = brackish-salt: 1000-10,000; S = salt: 10,000-20,000 and H = hyperhaline: $>20,000$ (**Table 1S**).

Alkalinity (in meq/L) determines the type and may vary from very low ($<0.5 \text{ meq L}^{-1}$) to extremely high ($>256 \text{ meq L}^{-1}$). The type includes a total of 10 “classes” of values coded from * to 9 (see **Table 2S**). In most natural waters with $4.5 \leq \text{pH} \leq 9.5$ (at 25°C , 1 bar), alkalinity is equal to the sum of HCO_3^- and CO_3^{2-} ([Stuyfzand, 1989](#)) or only HCO_3^- ion for $\text{pH} \leq$

8.3 (at 25°C, 1 bar), as is the case for the GTP waters. Nevertheless, it must be kept in mind that alkalinity is defined as the sum of all the bases dissolved in the aqueous solution and, therefore, it receives the contributions of several other dissolved anions (e.g., phosphate anions, namely PO_4^{3-} and HPO_4^{2-} entirely and H_2PO_4^- in part, OH^- , H_3SiO_4^- , H_2BO_3^- , HS^- , CH_3COO^- and other organic acid anions) and neutral species (e.g., NH_3), which may become even more important than HCO_3^- and CO_3^{2-} for high-pH Ca(Mg)-OH waters (e.g., [Bruni et al., 2002](#); [Cipolli et al., 2004](#)), formation waters ([Palandri and Reed, 2001](#)), and geothermal brines affected by boiling ([Cioni and Marini, 2020](#)).

The identification of subtypes is based on the dominant cations and anions, by using the triangular diagrams of $(\text{Ca}^{2+}+\text{Mg}^{2+})-(\text{Na}^++\text{K}^++\text{NH}_4^+)-(\text{Al}^{3+}+\text{H}^++\text{Fe}^{2+}+\text{Mn}^{2+})$ and $(\text{HCO}_3^-+\text{CO}_3^{2-})-(\text{Cl}^-)-(\text{SO}_4^{2-}+\text{NO}_3^-+\text{NO}_2^-)$ both prepared adopting the concentrations in meq L⁻¹.

Finally, BEX defines the *class* and identifies the cation exchange reactions occurring during the infiltration of either saline water in a fresh-water aquifer or freshwater into a saline/brackish aquifer containing several exchangers (e.g., including clay minerals, organic matter, and metal oxyhydroxides), as typically found in coastal areas. Indeed, during marine intrusion, a salinity increase affects the aquifer, and the following exchange reaction takes place:



where X indicates the cation exchanger. In the reaction, Na^+ is taken up by the exchanger, while Ca^{2+} is released to the aqueous solution. Since no reaction involves Cl^- , water composition changes from Na-Cl to Ca-Cl.

During the freshening processes, the reverse reaction of (1), induced by freshwater recharge, takes place. The cation exchanger adsorbs Ca^{2+} while Na^+ is released to the aqueous solution with the consequent evolution to the Na- HCO_3 type:



Based on these premises, the analytical cation concentrations of groundwater are commonly used through BEX to identify salinization or freshening tendencies occurring in coastal areas. Several base exchange indices were proposed by different authors as reported by [Stuyfzand \(2008\)](#) who recommended the use of the following:

$$\text{BEX} = (\text{Na}+\text{K}+\text{Mg}) - 1.0716 \text{ Cl (meq/L)} \quad (3)$$

where 1.0716 is the (Na+K+Mg)/Cl ratio in mean ocean water. Thus, the BEX index indicates how much the sum of Na, K, and Mg deviates with respect to the seawater contribution. Nevertheless, such a deviation from the marine cation composition may also be caused by other processes such as mineral dissolution/precipitation reactions ([Stuyfzand, 2008](#)). Owing to these uncertainties concerning the BEX index, we decided to ignore the BEX classes and to adopt a simplified code based only on the main type, type, and subtype. The simplified Stuyfzand's classification was used for the straightforward visualization of space trends on cross-sections and maps thanks to the effective and compact coding of the different water facies ([Figure 1a](#)). Finally, the methods of spatial interpolation Inverse Distance Weighting (IDW) and the ordinary kriging, obtained using suitable semi-variogram models to describe the spatial dependence of each parameter, were applied for the generation of iso-concentration maps.

4. Results and Discussion

4.1. Physical-chemical parameters

324 The main statistical indices of the physicochemical groundwater parameters of HU3 and HU1 aquifers are summarized
 325 in [Table 3S](#). The GTP groundwaters show a wide range of electrical conductivity (from 228 to 10,520 $\mu\text{S}/\text{cm}$) indicating
 326 that they range from very fresh to very saline, according to [Phocaides, \(2000\)](#). The pH from both aquifers exhibits a great
 327 variability from north to south of the plain, with values ranging between 5.9 and 8.9. The temperature range is from 14 to
 328 25.8 °C while Eh varies from -250 mV to 300 mV, the negative values mostly related to the HU1 aquifer although,
 329 somewhere, Eh values down to -106 mV were also measured in the HU3 groundwaters.
 330 From a general point of view, the main anomalies of the physicochemical parameters are recognized in the northern
 331 boundary of the plain (Sector B), along the NGFZ_ NW-SE_S, one of the main fault zones dissecting the study area.
 332 Waters in this sector show high values of conductivity, pH, and temperature, with pH values > 8 and a peak temperature
 333 of 26 °C. Furthermore, the northern sector is characterized by the most negative Eh values when compared to the other
 334 areas. The waters from the southern portion of the plain (sector F) are similar to those of the central sectors (C/D), having
 335 low conductivity (< 1000 $\mu\text{S}/\text{cm}$), positive Eh values and relatively low temperatures (< 18 °C). Only one restricted area,
 336 along the coastline (Sector E) and confined by the Palmi high (Scinà locality), revealed anomalous conductivity values,
 337 > 1500 $\mu\text{S}/\text{cm}$.
 338 A significantly different spatial distribution of the physicochemical parameters is shown by the two main aquifers (HU3
 339 and HU1). The 21 samples of deep artesian wells of HU1 (red dots in [Figure 1a and 1Sa](#)) show a spatial heterogeneity
 340 much greater than that observed for HU3. The HU1 groundwater shows the highest temperature (mean of 18.6 ± 3.2 °C),
 341 highest conductivity (mean of 1224 $\mu\text{S}/\text{cm}$ with peaks up to 10,520 $\mu\text{S}/\text{cm}$), and pH (values > 7.5) as well as negative Eh
 342 values, with the main anomalies found mainly in proximity of the northern boundary of the plain (Sector B). The HU3
 343 groundwater shows mean temperature of 17.4 ± 1.3 °C, variable conductivity values (between 249 and 4049 $\mu\text{S}/\text{cm}$), and
 344 pH between 5.9 and 8.5 as well as, in most cases, positive Eh values. In the northern portion of the plain (northward to
 345 the Mesima mouth), groundwater of HU3 show relatively high conductivity ($\text{EC} > 1000$ $\mu\text{S}/\text{cm}$) and localized negative
 346 Eh values. Nevertheless, the main anomaly ($\text{EC} > 1500$ $\mu\text{S}/\text{cm}$) in HU3 occurs in the south-west (sector E), close to Scinà
 347 ([Figure 1a](#)). Finally, the middle and inner portions of the plain (around Rizziconi, Taurianova and Polistena villages),
 348 display the lowest temperature and electrical conductivity values.
 349 The physicochemical parameters of main rivers (Budello, Mesima, Metramo and Petrace Rivers) are in most cases similar
 350 to those shown by HU3 groundwater. River waters have mean temperature of 15.1 ± 2.9 °C, mean electrical conductivity
 351 of 393 ± 79 $\mu\text{S}/\text{cm}$, pH between 7.2 and 8.7 and positive Eh values. Regarding the recharge area, the Capo Vaticano High
 352 (PG16, PG17, SG4 and PG71), Palmi High (PG94) and Aspromonte/Serre Massif (PG36) groundwaters differ from the
 353 HU1, HU3 groundwater and surface waters due to the distinct lithological-structural framework. These samples were
 354 collected close to the GTP outer boundaries, where the groundwater may have undergone a water-rock interaction extent
 355 higher with respect to the rainwater infiltrating at high altitudes. Waters from the Aspromonte/Serre Massif show the
 356 lowest temperature (14.5 °C) and electrical conductivity (364 $\mu\text{S}/\text{cm}$) values and have relatively high pH (7.9), whereas
 357 waters from the Capo Vaticano High (San Giovanni Basin) have electrical conductivities up to 1068 $\mu\text{S}/\text{cm}$ and pH
 358 between 7.1 and 7.3. Finally, the Palmi High (southern sector) water has an electrical conductivity of 655 $\mu\text{S}/\text{cm}$ and a
 359 pH of 6.8, i.e., the lowest one among the three recharge areas.


361 **4.2. Water classification**

362 According to the Durov plot ([Figure 2](#)), it is possible to distinguish the different geochemical facies that characterize the
 363 groundwaters (from HU1, HU3, and recharge area) and surface water.

364 (*i - shallow aquifer*). Shallow groundwaters (HU3) are ascribable to five main chemical facies, namely: Ca-HCO₃ (55
 365 samples), Na-HCO₃ (6 samples), Ca-SO₄ (8 samples), Na-Cl (7 samples) and Ca-Cl (2 samples) (Table 1S). Groundwater
 366 samples, showing comparable composition with samples of recharge areas (orange stars in Figure 2), are mainly Ca-
 367 HCO₃ since Ca largely prevails over Mg, with a Ca/Mg ratio (in meq L⁻¹) of 3.3 ± 1.4. These chemical characteristics are
 368 probably acquired through water-rock interaction between meteoric waters and Ca-rich carbonate minerals, such as calcite
 369 and dolomite (Appelo and Postma, 2005; Apollaro et al., 2009, 2013; White et al., 1999), whose dissolution rate is several
 370 orders of magnitude higher than silicate rocks (e.g., Marini, 2006 and reference therein). Calcite is present as an accessory
 371 phase in granitoid rocks with concentrations varying between 20 and 18,800 mg/kg (average value of 2520 mg/kg; White
 372 et al., 1999, 2005). Thus, meteoric waters of the study area rapidly acquire the Ca-HCO₃ composition through dissolution
 373 of disseminated calcite present in the crystalline basement (along the recharge areas) and in the sedimentary rocks derived
 374 from the erosion of the crystalline basement. The process is indicated by the following reaction:



378 An ideal evolution of this process would involve the distribution of points along a horizontal line in Figure 3a. Instead,
 379 Ca-HCO₃ waters also show a gradual increase in Na⁺ and K⁺ related to the possible concomitance of secondary processes
 380 such as the interaction with crystalline rocks, anthropogenic inputs, and ionic exchange reactions but it is difficult to
 381 understand which of these processes explain(s) the enrichment in alkali ions of Ca-HCO₃ waters. Ionic exchange reactions
 382 (1) and (2) are described by iso-salinity trends and account for the iso-salinity evolution from the Ca-HCO₃ to the Na-
 383 HCO₃ facies.



384
 385 **Figure 2** - Durov diagram showing the GTP water facies. Differences from the Ca-HCO₃ type are highlighted by colour shading,
 386 namely: yellow for Na-HCO₃ of sectors B, C, D and F; orange for Na-HCO₃(Cl) to Na-Cl of sector B; lilac for Na-Cl waters of sector
 387 E (Scinà area); and purple for Ca-SO₄ and Ca-Cl waters of sector C. FW and SW represent freshwater and seawater, respectively.


389 The Ca-HCO₃ waters from HU3 are mainly found in the eastern portion of the plain (Sector *A* – [Figure 1Sb](#)) but they are
 390 the dominant type of the shallow aquifer (70%). This type shows low but variable electrical conductivity, between 249
 391 and 1622 µS/cm (from oligohaline [g] to fresh-brackish [f] waters), with the highest value recognized in proximity of the
 392 GTP northern boundary (Sector *B*).

393 The north-central portion of the GTP (Sector *C*), where the inland field of dunes crops out, is characterized by the presence
 394 of Ca-SO₄ waters. This chemical type has relatively low conductivity, between 400 and 825 µS/cm (Fresh [F]) and
 395 exclusively pertain to HU3. Given the absence of gypsum- or anhydrite-bearing evaporites in the study area, this chemical
 396 type could be linked to local anthropic factors or oxidative pyrite dissolution ([Rickard, 2012; Rickard et al., 2017](#))
 397 followed by calcite dissolution (the processes controlling the development of the different chemical types will be
 398 discussed in depth in the next section).

399 The Na-HCO₃ waters from HU3 occur in the northern (Sector *B*, between 0.9 and 1.6 km away from the coastline) and
 400 south-central parts (Sector *D*, between 1 and 9 km away from the coastline) of GTP close to Nicotera Marina and Budello
 401 River, respectively ([Figure 1a](#)). In the northern sector, the Na-HCO₃ samples from HU3 (PG47 and PG75) reflect the
 402 composition and the relatively high salinity recognized in the local recharge water. In fact, waters from the Capo Vaticano
 403 High (e.g., PG17 representative of the San Giovanni Basin) show high conductivity values when compared with those
 404 from the eastern (Aspromonte/Serre) and southern (Palmi High) recharge areas, respectively. On the other hand, for the
 405 Na-HCO₃ waters from HU3, collected close to Budello River (Sector *D*), a distinction between low and relatively high
 406 conductivity (from 299 up to 835 µS/cm, respectively) waters is to be made. The low-salinity waters show high Ca/F
 407 molar ratio (up to 27) and are undersaturated in calcite. These characteristics are typical of immature waters deriving from
 408 the Na-Cl rain waters which are typically found in coastal areas and can be considered strongly diluted seawater ([Appelo
 409 and Postma 2005](#)). The evolution from this meteoric composition (very low-TIS G-Na-Cl) to the low-TIS G/g-Na-HCO₃
 410 facies occurs through limited water/rock interaction with the upstream crystalline-metamorphic successions ([Fuoco et al.,
 411 2022](#)). In contrast, the higher salinity samples show oversaturation with respect to calcite and a lower Ca/F molar ratio
 412 (11.5), suggesting a prolonged water/rock interaction with crystalline-metamorphic rocks ([Fuoco et al., 2022; Apollaro
 413 et al., 2015](#)). Consequently, the overall geochemical evolution can be schematized as: very low TIS G-Na-Cl → low-TIS
 414 G/g-Na-HCO₃ → g/F-Ca-HCO₃ → relatively high-TIS F-Na-HCO₃ ([Fuoco et al., 2022](#)). However, the occurrence of
 415 freshening processes cannot be excluded since during direct cation exchange, calcium concentration decreases favouring
 416 the dissolution of F-rich phases (e.g., fluorite) or F-bearing minerals (e.g., apatite, muscovite, and amphibole), leading to
 417 a low Ca/F molar ratio ([Chen et al., 2020](#)).

418 The Ca-Cl composition was only identified in two wells from Nicotera Marina (PG74) and close to Rizziconi (PG99).
 419 Generally, the chemical Ca-Cl type is mostly due to marine ingression and subsequent cation exchange, according to
 420 reaction (1). Nevertheless, both samples represent isolated cases located at different distances from the coast (2 km and
 421 10 km, respectively). For the former an anthropogenic source can be hypothesized. Sample PG74 is indeed localized in
 422 proximity of a water treatment plant for drinking water purposes and could account for the anomalous release of Na-Cl
 423 waters into the HU3 aquifer. Furthermore, other wells located between PG74, and the coastline typically show a Ca-HCO₃
 424 compositions. Regarding sample PG99, the distance from the coast (10 km) allows to exclude any ion exchange processes
 425 linked to seawater intrusion. In this sample, the low Cl/Br molar ratio confirms anthropogenic contamination (see *section*
 426 *4c*). Finally, the Na-Cl group (samples PG38, PG66, PG67, PG68, PG69 and PG70) is mainly found in the southern
 427 coastline (Sector *E*), in proximity of Scinà. These samples, as highlighted in [Figures 3a and 3b](#), show the highest
 428 conductivity values, between 1144 and 4049 µS/cm (from Fresh [F] to brackish-salt [b]), in the shallow aquifer. In


429 particular, samples PG38, PG66, PG67, and PG68 could be produced through either seawater addition to low-salinity Ca-
 430 HCO_3 waters or ionic exchange according to reaction (2) involving Ca- HCO_3 waters of higher salinity, as indicated by
 431 lines c, d and a, b, respectively (Figure 3a). Nevertheless, the chemical composition of PG69 and PG70 is unlikely to be
 432 related to ionic exchange as it would have involved Ca- HCO_3 waters characterized by very high salinities, much greater
 433 than the observed values. Since these two samples lie close the seawater dilution line (Figure 3b - data from Drever,
 434 1988), it is possible to suggest the presence of direct seawater intrusion, though at a relatively low degree.
 435




436 **Figure 3** - Correlation diagram of $(\text{Ca}^{2+} + \text{Mg}^{2+})$ vs. $(\text{Na}^+ + \text{K}^+)$ for (a) low and (b) high salinity samples from the GTP waters. Iso-salinity
 437 lines are drawn for reference. Lines a, b, c, and d are discussed in the text. (c) Activity diagram (sensu Helgeson 1968) showing the
 438 stability fields of low-albite, kaolinite, paragonite, gibbsite, Na-beidellite and saturation lines with respect to chalcedony and opal.
 439 Binary plots (d) $\text{Ca}^{2+} + \text{Mg}^{2+}$ vs. $\text{HCO}_3^- + \text{SO}_4^{2-}$, (e) Cl^- vs. Na^+ and (f) EC vs. Na/Cl showing the water samples from the Gioia Tauro
 440 Area.
 441
 442

443 (ii - deep aquifer). Deep groundwater (HU1) (represented by diamonds of different colour in Figure 2) can be divided
 444 into three distinct groups, as follows: Ca- HCO_3 (7 samples), Na- HCO_3 (10 samples), and Na-Cl (4 samples) (Table 1S).
 445 The Ca- HCO_3 waters, with a Ca/Mg ratio (in meq/L) of 1.82 ± 0.27 , show very low salinity (oligohaline waters [g]) and
 446 are mainly found along the southern and eastern boundary of the plain, that is, in sectors F and A. The Na- HCO_3 waters
 447 have low concentrations of Cl^- (<105 mg/L), Ca^{2+} (<45 mg/L), and Mg^{2+} (<30 mg/L). This group of waters represents the
 448 dominant facies of HU1 (48%) characterized by intermediate conductivity, between 350 and 878 $\mu\text{S/cm}$ (from oligohaline
 449 [g] to fresh [F] waters) and is well distributed along the southern, central and northern portions of the study area (Sectors
 450 B, C, D and F). Freshening processes generally control this facies in coastal areas, mainly for the cation exchange reaction
 451 (2) (Appelo and Postma, 2005). However, the Na- HCO_3 waters with low salinity can be produced by relatively prolonged
 452 interaction between meteoric water and sedimentary rocks deriving by the dismantle of crystalline rocks, governed by the

conversion of CO_2 to HCO_3^- in confined aquifers where the residence time of water is comparatively long. The process is schematically represented by the incongruent dissolution of albite, accompanied by precipitation of kaolinite and possibly a silica mineral (Garrels, 1968):



If albite dissolution occurs at undersaturation conditions with calcite, the concentrations of Na^+ and HCO_3^- are expected to increase in a 1:1 ratio, as dictated by reaction (5). In contrast, if albite dissolution is accompanied by calcite precipitation, then the following reaction would take place:



Reaction (6) implies constancy of HCO_3^- concentration and the $(\text{Ca}^{2+} + \text{Na}^{2+})$ concentrations in meq L^{-1} . This means that the process is undistinguishable from direct cation exchange, that is, reaction (2). If so, ionic exchange and relatively prolonged dissolution of silicate minerals would virtually produce overlapping effects that are difficult to be discerned. Nevertheless, rock dissolution requires participation of low-salinity Ca- HCO_3 waters (in line with the available evidence), whereas ionic exchange necessitates the involvement of Ca- HCO_3 waters of progressively higher salinity, which were not found in HU1. As shown in Table 1S, almost all Na- HCO_3 waters are in equilibrium or slightly oversaturated with respect to calcite. This makes it further complex to distinguish between the two processes. To provide further details, the decimal logarithms of the Na^+/H^+ activity ratio and the decimal logarithm of SiO_2 activity are plotted into the activity diagram of Figure 3c. This stability diagram was preferred to relevant saturation indices because analytical Al concentrations may be affected by incomplete removal of suspended colloidal particles and thermodynamic data on Al-bearing solid phases are controversial (e.g., Cioni and Marini 2020). The HU1 Na- HCO_3 waters fall in the low-albite stability field or close to the boundary separating the kaolinite and low-albite stability fields, thus indicating a greater evolution degree compared to the Ca- HCO_3 waters and the possible occurrence of reaction (6) at saturation state with calcite. Although this evidence is not conclusive, if the $\text{Ca}^{2+} - \text{Na}^+$ exchange were controlled by freshening there would be no reason to find water samples close to the kaolinite/low-albite boundary. The origin of this specific chemical type (Na- HCO_3 waters) will be debated in detail in the next section.

Finally, the Na-Cl waters are found in both the outer northern and the southern portions of the plain (Sector B and F) and have the highest conductivity values: from 1858 to 10520 $\mu\text{S}/\text{cm}$ (from fresh-brackish [f] to brackish-salt [b] waters). Most Na-Cl HU1 groundwaters have Cl/SO_4 ratio (meq L^{-1}) of 2 ± 0.7 , with the exception of sample PG81 (sampled by a well 436 m depth) whose Cl/SO_4 ratio is ~ 550 , possibly due to the occurrence of bacterial sulphate reduction (BSR) under strongly reducing conditions (Marini et al., 2000), as supported by the Eh value (-194 mV). Additionally, as shown in Figure 3b, the Na-Cl HU1 groundwater is found somewhat to the left of the seawater Na-Cl dilution line of shallow groundwater, suggesting that the chemical groundwater evolution is different in the two aquifers.

(iii – river water). Surface water of the Budello, Metramo, Mesima and Petrace rivers belong mainly to the Ca- HCO_3 type. This type is typical for freshwater recharge from the northern, southern, and eastern part of the study area. The presence of an isolated Na- HCO_3 water near the Budello river mouth (sample FG1) is probably due to freshening.

(iv – recharge water). Most samples collected in proximity of major recharge areas have Ca- HCO_3 composition and relatively low salinity (g for the Aspromonte/Serre Massifs [PG36] and F for the northern and southern areas). The only exceptions are samples PG17 and PG94, representative of the San Giovanni Basin and Palmi High, respectively, which are Na- HCO_3 .

Overall, the Durov square diagram (Figure 2) reveals that most waters from HU1 and HU3 are positioned in the mixing area connecting the typical (extreme) Ca-HCO₃, NaCl (seawater) and Ca(Mg)-HCO₃ waters, with little or no shifting caused by secondary processes (e.g., direct and inverse cation exchange or prolonged dissolution of silicate minerals). The right and left corner areas in the graph are usually considered representative of fresh water (FW) and seawater (SW), respectively (e.g., Giménez-Forcada 2010; Giménez-Forcada and Sánchez San Román 2015). Nevertheless, the rain waters in coastal areas are strongly diluted seawater (Appelo and Postma 2005) and, therefore, they are indistinguishable from seawater in this plot. Since meteoric waters evolve rapidly from initial marine-atmospheric Na-Cl compositions towards Ca-HCO₃ compositions through quick calcite dissolution (see above), sample points move from SW towards FW in the Durov square diagram and not vice versa. Consistent with these expectations, the low-salinity Ca-HCO₃ waters of the GTP are found close to the centre of the Durov square plot, whereas most other Ca-HCO₃ waters are approaching the upper right corner of this plot as salinity increases.

In HU1, 48% of samples are Na-HCO₃ and positioned along a nearly vertical alignment representative of direct cation exchange or dissolution of Na-Al-silicates and Na-silicates. The Ca-HCO₃ type, recognized in HU1, is characteristic of the eastern and southern boundaries, in proximity of the main recharge areas. These samples, as shown in Figure 2, fall close to the diagonal line connecting the upper right and the lower left corner areas but they are slightly shifted downward due to either direct cation exchange or dissolution of Na-Al-silicates and Na-silicates. The HU1 waters sampled in proximity of the northern boundary (Sector B) fall between the fields (i) and (c) of the Durov square diagram. These waters show relatively high temperature (peak of 26 °C). Their composition is probably conditioned by the NGFZ_NW-SE_S, that favours the uprise of deep Na-Cl thermal waters of high salinity, similar to sample PG81 (Sector B), locally intruding the HU1. In the shallow aquifer (HU3), salinization occurring in the southern portion (Scinà Area – Sector E) is probably caused by lateral seawater flow from the west. As shown in Figure 2, the waters samples from the Scinà area fall in proximity of the conservative mixing zone between freshwater and seawater or slightly below it. In this sector, the groundwaters are affected by saltwater intrusion but do not seem to be significantly influenced by inverse cation exchange reaction ($\text{Na}^+ \rightarrow \text{Ca}^{2+}$) as indicated by the absence of Ca-Cl type waters (see the next section). As reported in section 2, Scinà is characterized by a peculiar hydrogeological setting where the shallow aquifer, made by the fluvio-deltaic sediments of the Petrace River and coastal deposits, lies directly in contact with the metamorphic basement. In this framework, the exchangers, necessary for the occurrence of inverse cation exchange and the production of Ca-Cl groundwaters, are expected to be present in small amounts or absent.

523

524 4.3. Origin of solutes

The processes controlling the development of the different chemical types is examined by using binary diagrams. Although binary plots involving mobile and conservative species give a partial view of the relationships among the water samples, they can still be useful to verify whether the spread of sample points is consistent with the trends expected for different processes probably occurring during water circulation. In the HU1 and HU3 aquifers relevant processes seem freshening, dissolution of silicate minerals, bacterial sulphate reduction and mineral precipitation. In particular, given the geological framework of the study area, the B, F⁻ and SiO₂ variability, potentially discriminating species of water-rock interaction with crystalline rocks (Fuoco et al., 2022), was considered. These elements are normally compared with chloride that can be considered a conservative tracer of water evolution (Flury and Prapitz, 1993). Furthermore, chloride concentrations in seawater (ca. 545 meq/L, Drever, 1988) are significantly higher than those expected in fresh unpolluted groundwater. In this sense, the chloride concentration in the GTP aquifers may be related to either dissolution of Cl-bearing minerals from the aquifer host rocks, mixing with seawater at different degrees or, chloride influx with recharge

water (marine aerosols rich in chloride due to sea proximity). Since no Cl-bearing minerals are reported in the reservoir, the climate conditions of the study area (evaporation), connate waters, direct seawater intrusion, and/or the marine aerosols can be identified as the main causes of increasing chloride concentrations.

From a lithological point of view, the GTP basin is filled by siliciclastic and carbonate successions. In this lithological framework, a balance among Ca^{2+} , Mg^{2+} , HCO_3^- and SO_4^{2-} concentrations is expected whenever these solutes are released from dissolution of calcite, dolomite and gypsum (Sarker et al., 2021) and the correlation between $(\text{Ca}^{2+} + \text{Mg}^{2+})$ and $(\text{HCO}_3^- + \text{SO}_4^{2-})$ is expected to be close to the 1:1 ratio line (Khairy and Janardhana, 2013; Nasher et al., 2013; Fadili et al., 2015; Sarker et al., 2021). The graphical plot (Figure 3d) of $(\text{Ca}^{2+} + \text{Mg}^{2+})^+$ vs. $(\text{HCO}_3^- + \text{SO}_4^{2-})$ shows that most GTP waters belonging to the Ca- HCO_3 , Ca- SO_4 , Na-Cl, and Ca-Cl water types fall along the 1:1 line. As observed, no sample exhibits Ca^{2+} and Mg^{2+} excess testifying the absence of inverse cation exchange. On the other hand, most Na- HCO_3 waters from HU1 and HU3 are located below the 1:1 line because their HCO_3^- concentration is chiefly balanced by Na. This could be due to either direct cation exchange process or occurrence of silicate weathering reactions that also produce an increase of dissolved Na^+ and K^+ .

In the Na^+ vs. Cl^- plot (Figure 3e) most waters fall in proximity of the 1:1 halite dissolution line, apart most Na- HCO_3 waters from HU1 and HU3 that are marked by a Na^+ -excess. These waters being positioned above the halite dissolution and seawater dilution lines because their Na^+ concentration is chiefly balanced by HCO_3^- . The departure of the Na- HCO_3 waters from the halite and seawater dilution lines is even more evident in the plot of the Na/Cl ratio vs. EC ($\mu\text{S}/\text{cm}$) (Figure 3f) in which most samples fall close to the halite dissolution and seawater dilution lines, indicating a direct mixing with seawater in controlling the Na/Cl ratio.

The HU3- and HU1-NaCl waters have different Na/Cl ratios (Figure 3e). While HU3 NaCl groundwater falls across the seawater dilution line, testifying a direct seawater intrusion, the HU1 NaCl samples are positioned above the seawater dilution line, highlighting ratios comparable to those recognized in the Galatro thermal groundwater. Only sample PG81 (B-NaCl, sector B), in HU1, falls along the 1:1 Na/Cl line. The location (5 km far from the coast) and depth (> 430 m) of PG81 and the absence of evaporitic successions (halite) in the GTP suggest a possible origin linked to a mixing with old connate waters entrapped during the marine transgression.

The HU1- and HU3-Na HCO_3 waters show a positive shift of the Na/Cl ratio at relatively low and comparable salinity, testifying the presence of secondary processes increasing Na^+ concentration. With the aim to improve the knowledge about the main processes able to explain the evolution towards Na- HCO_3 compositions, and the origin of the brackish waters sampled in the northern side (sector B), mass balance calculations were carried out on the PG81 brackish water. The $\text{Na}_{\text{marine-derived}}$ concentration, i.e., the concentration expected if mixing with seawater were the only process able of providing Na^+ to solution, was calculated considering the $\text{Cl}_{\text{measured}}$ concentrations (obtained in this study) and those of Cl_{SW} and Na_{SW} representative of average seawater (Drever, 1988):

$$C_{i, \text{ marine-derived}} = (C_{i, \text{ SW}}/\text{Cl}_{\text{SW}}) * \text{Cl}_{\text{measured}}. \quad (7)$$

where C_i is the concentrations of i-th component.

The $\text{Cl}_{\text{measured}}/\text{Na}_{\text{marine-derived}}$ ratio ($92.97/79.77 = 1.165$ in equivalent unit) is consistent with the mean seawater ratio (1.166 in equivalent unit) as shown in Figure 3e. Therefore, it can be inferred that direct exchange processes do not take place in the brackish water PG81. Thus, the increase in Na^+ is likely due to water-rock interaction processes with the rocks of the crystalline units. Accordingly, the Na concentration derived through dissolution of silicate minerals, Na_{WR} , was calculated as the difference between the analytical value ($\text{Na}_{\text{measured}}$) and $\text{Na}_{\text{marine-derived}}$. The Na_{WR} turned out to be 12.39

577 meq L⁻¹, which corresponds to the length of the red arrow in [Figure 3e](#), whereas the corresponding Ca_{WR} (difference
578 between Ca_{measured} and Ca_{marine-derived}) resulted to be -2.59 meq L⁻¹ which is obviously meaningless because concentrations
579 cannot be negative. The negative Ca_{WR} value is probably due to the occurrence of calcite precipitation before sampling.
580 Accepting this explanation, the initial (before calcite precipitation) Ca and HCO₃ concentration are at least 3.49 and 21.14
581 meq L⁻¹ (for Ca_{WR} = 0) instead of the analytical counterparts (0.9 and 18.55 meq L⁻¹, respectively). In the correlation
582 diagram of [Figure 4a](#):

- 583 (i) the HU1-Ca-HCO₃ and HU3-Ca-HCO₃ waters are distributed along a horizontal trend, as expected based on reaction
584 (4);
585 (ii) the HU1-NaHCO₃ and HU1-NaCl waters (including sample PG81 corrected for the effect of calcite precipitation) are
586 situated along a vertical trend and the HU3-Na-HCO₃ waters occupy intermediate positions consistent with the acquisition
587 of Na⁺ and the corresponding loss of Ca²⁺ as exemplified by reaction (6).

588 Based on the above considerations and given the distribution of the Na-HCO₃ waters in both HU1 and HU3, even found
589 at distances of several km from the coast, we can conclude that most Na-HCO₃ waters acquire alkaline elements (chiefly
590 Na⁺) by prolonged water-rock interaction with crystalline rocks and/or sediments derived from their erosion. As already
591 mentioned, the evolution of this process is represented by lines of slope -1 in [Figure 4a](#), which are representative of a Ca-
592 Na exchange process at constant salinity. In many natural systems, after achieving the saturation condition with respect
593 to carbonate phases (primarily calcite), the dissolution of silicates (mainly feldspars and micas), governed by the
594 conversion of CO₂ to HCO₃⁻, indeed occurs with calcite precipitation and results in the transition from the Ca-HCO₃ to
595 the Na-HCO₃ facies at nearly constant salinity (e.g., [Pastorelli et al., 2001](#) and reference therein).

596 However, the F⁻ vs. SiO₂ diagram ([Figure 4b](#)) indicates how the discrimination of pure water-rock interaction with
597 crystalline rocks from ion-exchange processes is far from simple to be recognized. It should be noted that fluoride is likely
598 to have a quasi-conservative behaviour in the waters under investigation, since all samples are undersaturated with respect
599 to fluorite ([Table 1S](#)). The diagram of [Figure 4b](#) suggests that Na-HCO₃ samples are controlled by the three distinct
600 endmembers A, B, and C, whose characteristics are as follows:

- 601 • Endmember A has high F (~1.7 mg/L) and low SiO₂ (~ 6 mg/L) and is probably related to direct ion exchange,
602 which favours the decrease in calcium, thereby increasing the solubility of fluorine-rich phases ([Chen et al.,](#)
603 [2020](#)).
- 604 • Endmember B has high F (~2 mg/L) and high SiO₂ (~45 mg/L) and is probably related to direct water-rock
605 interaction with crystalline rocks ([Fuoco et al., 2022](#)). Furthermore, the occurrence of volcanoclastic pumice-
606 rich deposits was reported in GTP ([De Rosa et al., 2008](#)). These lithotypes typically contain F-bearing minerals
607 acting as F-sources ([Dharmagunawardhane et al., 2016](#)). Moreover, the samples richer in endmember B have the
608 highest boron concentrations ([Figure 4c](#)) which are also typical of waters interacting with volcanoclastic deposits
609 ([Dharmagunawardhane et al., 2016](#)).
- 610 • Endmember C has slightly high SiO₂ (~37 mg/L) and very low F and is probably related to limited water-rock
611 interaction processes with crystalline rocks or mixing with Ca-HCO₃ and SiO₂-rich waters.

612




Figure 4 – Binary plots (a) Ca_w vs. Na_w , (b) SiO_2 vs. F , (c) B vs. F , (d) HCO_3^- vs. Ca , (e) HCO_3^- vs. Ca/Na and (f) Ca vs. SO_4 showing the water samples from the Gioia Tauro Area. BSR: Bacterial Sulphate Reduction. PG81CCP: corrected for calcite precipitation.

The different F/SiO_2 ratios are possibly suggesting the presence of different recharge areas. If, on one hand, the evidence confirms the presence of a water-rock interaction process with crystalline units and/or volcanoclastic deposits (high F , B and SiO_2 concentration and Na-HCO_3 waters recognized in the recharge areas), on the other hand, it is not justified to exclude a priori the presence of the cation exchange process, especially in those samples showing low SiO_2 concentrations. In the diagrams of Figures 4d and 4e, some Ca-HCO_3 samples from HU3 show a positive HCO_3^- shift with respect to the 1:1 and local mixing lines. These samples are localized in the northern portion of the study area (sector B) which is characterized by several peat levels (Cianflone et al., 2021). Furthermore, the waters have negative Eh values, typical of a reducing environment, which is favourable for the occurrence of the bacterially mediated sulphate reduction through the following reaction (Seal, 2006; Vespasiano et al., 2014):



where CH_2O is representative of organic matter (e.g., Berner and Berner, 1996; Stumm and Morgan, 1996; Fidelibus, 2003; Barker et al., 1998; De Montety et al., 2008), although the organic substances actually involved in this process might have a chemical composition other than CH_2O thus leading to a reaction stoichiometry different from that of reaction (8), as discussed by Marini et al., (2000). Irrespective of the stoichiometry of the BSR reaction(s) actually occurring, the BSR process determines a decrease of dissolved sulfate and a corresponding increase of sulfide species (which are usually removed from the aqueous solution, at least partly, through H_2S degassing and/or pyrite precipitation) as well as an increase of dissolved carbonate species (mainly HCO_3^- , in most cases), often triggering calcite precipitation and a consequent loss of dissolved Ca . In the SO_4^{2-} vs Ca^{2+} diagram (Figure 4f), the samples affected by BSR and possibly

637 calcite precipitation as well (situated in the grey area) are positioned below the local $\text{SO}_4^{2-}/\text{Ca}^{2+}$ evolution line confirming
638 the occurrence of these processes. Furthermore, the SO_4^{2-} versus Ca^{2+} plot shows that all water samples, both from the
639 shallow and deep aquifers, have $\text{SO}_4^{2-}/\text{Ca}^{2+}$ ratio < 1 , with the exception of the Ca- SO_4 waters from HU3, which are
640 slightly below the $\text{SO}_4^{2-}/\text{Ca}^{2+} = 1$ line. For these samples, gypsum dissolution was rejected since there is no evidence of
641 gypsum in the aquifer matrix and the sulfate vs. calcium plot indicated an evolution that could not be easily explained by
642 the dissolution of this phase, but it is more likely attributable to anthropogenic factors (see below).

644 4.4. Anthropogenic and deep hot systems impact on water chemistry

645 The Cl/Br ratio has successfully been used to distinguish between various sources of anthropogenic and naturally
646 occurring pollutants in groundwater (Alcala and Custodio, 2008; Chien and Lautz, 2018). Due to the conservative
647 behavior of Cl and Br ions, insights on the sources affecting the groundwater systems can be highlighted when peculiar
648 Cl/Br ratios are evidenced. The ocean has a constant molar Cl/Br ratio of 650 (Drever, 1988), which is slightly lower than
649 the local Tyrrhenian value (790 ± 90) recognized along the Sicilian coast (Deias unpublished data). Coastal rainfall shows
650 a similar Cl/Br ratio due to the Cl^- and Br^- input along the wind path from the ocean. Inland rainfall (especially in semi-
651 arid and arid areas) may have relatively lower Cl/Br ratios since Cl^- is removed by deposition of cyclic salts in coastal
652 areas (Davis et al., 2001; Edmunds, 2001). Halite dissolution would result in high Cl/Br ratios (up to 1000), as Br does
653 not enter the halite lattice during seawater evaporation (Cartwright et al., 2006). The interaction of groundwaters with Cl-
654 bearing contaminants such as domestic wastewater and fertilizers may also give a high Cl/Br signal, as well as the decay
655 of Cl-bearing organic compounds, while the degradation of Br-bearing organic substances may reduce Cl/Br ratios (Flury
656 and Prapitz, 1993).

657 Bromide concentrations in the GTP aquifers range from 0.0002 to 0.14 meq L^{-1} , and are significantly correlated with
658 chloride, with all the groundwater samples situated along the theoretical seawater mixing line or relatively close to it
659 (Figure 5a). The strong correlation observed between Br^- and Cl^- is less evident for those samples characterized by low
660 salinity, suggesting the occurrence of secondary processes or different anthropogenic inputs. As reported by Alcala and
661 Custodio (2008), anthropogenic processes can modify current or natural relic sources of salinity. In their work, the authors
662 collected several water samples affected by natural and anthropogenic sources and defined six groups of groundwater as
663 a function of the Cl/Br ratio: rainwater, unpolluted old phreatic water, recent phreatic water, polluted phreatic water,
664 saline and hypersaline relic seawater, and water affected by wastewater.

665 Most GTP samples fall in the “rainwater” group (blue circle in figure 5b), which is controlled by a prolonged dry period
666 during the summer and autumn followed by short, intense rainfall events and is characterized by $10 < \text{Cl} < 100 \text{ mg/L}$ and
667 Cl/Br ratios slightly below the seawater value. Line I in figure 5b is a horizontal evaporation line involving unpolluted
668 rainwater and producing uncontaminated old phreatic water having Cl/Br ratios between 600 and 700 (mol L^{-1}) for Cl
669 concentration between 10 and 100 mg/L. The range of Cl/Br ratios is comparable to that of seawater and may be explained
670 by decades of airborne marine aerosols from the adjacent coast. Line II is a horizontal mixing line that refers to unpolluted
671 old phreatic waters with salinity derived from unpolluted marine aerosols, and saline and hypersaline relic seawater
672 entrapped in sediments. The relic seawater shows a Cl/Br ratio approaching that of modern seawater. The gradual
673 contribution of unpolluted freshwater to saline and hypersaline relic seawater is concentrated around a horizontal mixing
674 line. Across the line II fall the waters affected by mixing with brackish waters (e.g., connate waters) and direct seawater
675 intrusion including the samples representative of sector E (Scinà). Line III is a horizontal evaporation line starting from
676 present polluted rainwater, with average Cl/Br ratios around 500 (mol L^{-1}), producing phreatic waters with similar Cl/Br
677 ratios. The use of this recent phreatic waters for irrigation only produces evapo-concentration, which does not affect the

original Cl/Br ratio of regional bulk deposition. These samples are representative of the GTP areas in which intensive agricultural activities largely occur. Line IV is representative of polluted phreatic waters impacted by Br-based pesticides, as methyl-bromide, e.g., used in sandy areas for soil fumigation in greenhouse. The Cl/Br ratios vary from 500 (mol L⁻¹), like average rainwater, up to values lower than 200 (mol L⁻¹). Most GTP samples and, in particular, the Ca-SO₄ waters of HU3 fall into this group confirming the possible anthropogenic contamination. Finally, the contribution of wastewater from domestic food preparation and cattle feeding increases the Cl/Br ratios up to 1000 (mol L⁻¹) (line V).




Figure 5 – Binary plots (a) Cl vs. Br, (b) Cl vs. Cl/Br, (c) $\delta^{18}\text{O}$ vs. Cl and (d) $\text{NO}_3(\text{wr})$ vs. $\text{SO}_4(\text{wr})$ showing the water samples from the Gioia Tauro Area. See the main text for figure details.

The role of the anthropogenic component is further investigated by comparing the values of $\delta^{18}\text{O}$ vs. Cl, NO_3/Cl vs. Cl and the calculated nitrate, calcium and sulphate excess in the GTP waters. Their excess represents the difference in concentration, $X_{(\text{wr})}$, between the observed values and those expected if mixing with seawater were the only process (equation 7). As previously stated, the contribution of seawater is derived from the conservative ion Cl^- (Appelo and Postma, 2005).

In Figure 5c, the two main recharge endmembers characterized by enriched (local coastal rainwater) and depleted (Serre/Aspromonte recharge area) $\delta^{18}\text{O}$ values, respectively, are reported. The mixing lines between brackish water and freshwater that cause a shift towards more enriched values in terms of Cl and $\delta^{18}\text{O}$ values, are also drawn. HU1 samples are localized in the sector representative of the Aspromonte/Serre massifs recharge, testifying the limited contribution of the local coastal rainfall, while the Ca-SO₄ samples of HU3 fall in proximity of the local rainwater. HU3 Ca-SO₄ samples, localized on the eastern side of the Gioia Tauro harbor, have shown high NO_3 and SO_4 concentration (figure 5d). In this

sector, agricultural activities, urban settlements and the complete absence of basic services, such as sewers or connections to aqueducts for domestic use, can favor the increase of these two constituents in solution. In fact, a strong correlation between $\text{NO}_3(\text{wr})$ vs. $\text{SO}_4(\text{wr})$ is typically detected in agricultural areas with no other significant salinization processes in act (Menció et al., 2016) because of the application of sulfur and nitrogen fertilizers such as ammonium carbonate $[(\text{NH}_4)_2\text{CO}_3]$ and sulfur-based fungicides (Li, et al., 2019). On the other hand, the second group of the HU3 Ca- SO_4 samples, representative of waters collected in the San Ferdinando Area (north side of the harbor), have shown significantly lower NO_3 concentration. However, in both areas, the anomalies are also attributable to the harbor and waste-to-energy power plant SO_x emissions which are scavenged by rainwater (Barberi et al., 2021). In fact, in maritime transportation, SO_x emissions mainly originate from the combustion of marine fossil fuels (e.g., Tassi et al., 2013), with SO_2 being the most predominant produced oxide. SO_x gases can react with oxygen in presence of NO_2 resulting in production of sulfuric acid, which can eventually produce acid rains. The HU3 Ca- SO_4 samples have the lowest pH values (mean value: 6.6, down to 5.9) among the GTP groundwaters, confirming for this chemical type an origin likely linked to anthropic issues, including harbor activities.

The NO_3 concentrations in the GTP waters reach values of 1.66 meq/L, with 8% of samples (belonging to HU3) exceeding the law limit values (0.81 meq/L) according to the D.Lgs. 152/2006 and the World Health Organization drinking water guidelines. The NO_3/Cl ratio vs. Cl plot (Torres-Martínez et al., 2021; Awaleh et al., 2022; Taussi et al., 2022), to distinguish whether NO_3 was derived from agricultural, soil, manure or sewage inputs is represented in Figure 6a.

The diagram clearly indicates that most HU3 samples align between the agricultural fertilizer and sewage endmember sources, confirming what was already highlighted with the Cl/Br ratio. The few HU1 samples, evidencing significant NO_3 concentration fall within the range of the organic soil. Of interest is the right-side of the diagram where the high salinity samples are located. In this area, the samples collected in the Scinà area are arranged along the seawater mixing line (seawater NO_3/Cl molar ratio of 0.0000086 by Drever et al., 1988) to confirm the ongoing seawater intrusion process.

Finally, among the most controversial samples taken from GTP, the Na-Cl waters from HU1 are characterized by relatively high temperature, up to 26 °C. These samples fall in sector B, i.e., close to NGFZ_ NW-SE_ S. To evaluate the origin of these groundwaters and relevant water-rock interaction processes, the $\text{Ca}(\text{wr})$ vs. $\text{SO}_4(\text{wr})$ values are investigated (Figure 6b) and the concentrations of B and Li are compared with the overall average composition of minerals and rocks (anhydrite, carbonate, halite, shale, basalt and granite from Turekian and Wedepohl, 1961; Stewart, 1963; Kushnir, 1980; Reimann and de Caritat, 1998) and seawater in Figure 6c (Giggenbach, 1991; Apollaro et al., 2016).

In Figure 6b, the HU1 Na-Cl waters fall close to the dilution line of the Galatro thermal site, which is located in the eastern part of the plain along NGFZ_ NW-SE_ S. Both thermal circuits issue from an enhanced permeability belt closely related to the presence of this regional tectonic line, as claimed by Pizzino et al., 2004. The above-reported trend, not observed in any other samples of the GPT, confirms a direct relationship between the HU1 Na-Cl and the Galatro thermal waters. Finally, the diagram of Figure 6c shows that B and Li of the GTP waters are chiefly acquired by leaching of carbonate rocks. Immature waters of recharge areas and waters affected by seawater intrusion (Na-Cl waters of HU3 from sector E) fall, as expected, in proximity of the local seawater B/Li ratio. The Na-Cl waters of HU1 are distributed along the dilution line between the Galatro thermal and PG81 waters, corroborating the inferences drawn from the $\text{Ca}(\text{wr})$ vs. $\text{SO}_4(\text{wr})$ diagram (Figure 6b) and not excluding the contribution of connate waters sampled in sector B.




Figure 6 – (a) NO_3^-/Cl molar ratios versus Cl concentration (in μmol) of the groundwater samples from Gioia Tauro Plain. In the diagram the representative fields of rain, agricultural, soil, sewage, and manure inputs, adapted from [Torres-Martínez et al., 2021](#), [Awaleh et al., 2022](#), [Taussi et al., 2022](#) are reported. Binary plots (b) $\text{Ca}_{(\text{wr})}$ vs. $\text{SO}_{4(\text{wr})}$ and (c) Li vs. B showing the water samples from the Gioia Tauro Area. See the main text for figure details.

4.5. Fraction of seawater and average recharge altitude estimation

The average infiltration altitude (H_i , m a.s.l.) for the groundwaters of the GTP was calculated by means of the following equation ([Vespasiano et al., 2015b](#)):

$$H_{i,18\text{O}} = -\frac{\delta^{18}\text{O} + 5.25}{0.00210} \quad (9)$$

Results were used to reconstruct the maps of [Figure 3S](#). The Inverse Distance Weighting (IDW) method of spatial interpolation, often used in Geographical Information System applications, is applied for the generation of this infiltration altitude map. In proximity of the Mesima River mouth and the upstream portion of the Budello River, anomalously high average infiltration altitudes are identified (>150 m), probably due to water exchanges between the rivers (locally affected by evaporation) and the adjacent portions of the shallow aquifer and/or the shallow and deep aquifers. These values were not considered in the reconstruction of the distribution maps. As shown in [Figure 3S](#), HU1 and HU3 have different infiltration altitudes calculated based on the $\delta^{18}\text{O}_{\text{GW}}$ value (see below). HU3 is characterized by a constant decrease in infiltration altitude, from ~ 350 to 0 m a.s.l. moving towards the coast, that is, from east to west, as represented by the iso-

infiltration line. These infiltration altitudes are likely to be influenced by local rains, i.e., from rainwater infiltrating directly into GTP.

The influence of local precipitation appears to be less important for HU1. In fact, the deep aquifer has virtually constant infiltration altitude values throughout the plain (410 ± 84 m a.s.l.), suggesting that the meteoric recharge is mainly related to the Aspromonte Massif at the contact with the sedimentary filling of the plain.

Average chloride content ($21,339 \pm 134$ mg/kg), $\delta^{18}\text{O}$ value ($+1.43 \pm 0.10$ ‰ V-SMOW) and $\delta^2\text{H}$ value ($+8.1 \pm 0.4$ ‰ V-SMOW) of local seawater have been assessed using the data available for the Ionian Sea (Stenni et al., 1995, Gat, 1996, Pierre 1999, Schmidt, 1999). Then, the fraction of seawater, x_{SW} , in each mixed groundwater can be computed by solving the chloride mass balance for x_{SW} :

$$x_{\text{SW}} = \frac{\text{Cl}_{\text{MW}} - \text{Cl}_{\text{GW}}}{\text{Cl}_{\text{SW}} - \text{Cl}_{\text{GW}}} \quad (10)$$

where the subscripts MW, GW and SW refer to mixed groundwater, groundwater unmixed with seawater ($\text{Cl} = 11.04$ mg/kg) and seawater, respectively. Then, the isotope balances were solved for δ_{GW} :

$$\delta_{\text{GW}} = \frac{\delta_{\text{MW}} - \delta_{\text{SW}} * x_{\text{SW}}}{1 - x_{\text{SW}}} \quad (11)$$

thus, obtaining the $\delta^{18}\text{O}_{\text{GW}}$ and $\delta^2\text{H}_{\text{GW}}$ values of the groundwaters unmixed with seawater (Table 1S). These corrected isotope values (black line in Figure 7a) define the local meteoric line, which diverges from the Southern Italy (Longinelli and Selmo, 2003; Giustini et al 2016) and Northern Calabria (Vespasiano et al., 2015b) meteoric lines. Local meteoric waters have much higher d-excess values (Table 1S), attributable to non-equilibrium (kinetic) evaporation at relatively low humidity.

The seawater fractions of local groundwater generally range from 0.1% to 0.2%, with peaks between 1% and 7% for the samples collected in sector E – Scinà Area (HU3) and about 15% for sample PG81, which is likely mixed with connate brackish water. As shown in the Cl vs. $\delta^{18}\text{O}$ plot (Figure 7b), the $\delta^{18}\text{O}$ values of groundwater span a range of 1.5‰ units, attributable to differences in the location and/or altitude of the recharge areas. Evaporative effects are evident for waters collected from wells of large diameter open to the atmosphere and exchanging with it (e.g., PG6).



789

794

795

796

797

798

799

800

802

803

804

805

806

807

808

809

810

811

813

814

815

816

817 (i) in HU3 inputs of N-rich sewage and agriculture-related contaminants and SO_x emissions in proximity of the harbour
818 represent the main processes favoring nitrate increases over almost the entire plain and the evolution towards the Ca-SO₄
819 composition. Furthermore, BSR processes induced by oxidation of organic matter in sector *B* and groundwater-seawater
820 mixing in sector *E* represent secondary and localized processes.

821 (ii) Na-HCO₃ waters of intermediate salinity are the dominant hydro-facies in HU1 with a percentage of 48%. Mass
822 balance calculations and the distribution of some major and minor constituents (SiO₂, F and B) allow one to identify at
823 least two possible origins for this chemical type: (a) the prolonged water-rock interaction with crystalline rocks and
824 sediments deriving from their erosion, also comprising volcanoclastic deposits and (b) cation exchange processes, for the
825 samples having low SiO₂ concentrations.

826 Connate waters are present at depth in the study area as indicated by sample PG81, which was collected from a 436-m-
827 deep well drilled in sector *B* and is characterized by Na-Cl composition and high concentrations of Cl (~3300 mg/kg) and
828 B (11 mg/kg), and relatively high Li concentration (0.13 mg/kg). Other two waters also coming from sector *B*, specifically
829 from HU1, samples PG10 and PG76, have Na-Cl composition, relatively high temperature (26 °C), and the same B/Cl
830 and Li/Cl ratios of sample PG81, but significantly lower Cl concentrations, 647 and 358 mg/kg, respectively, due to a
831 larger extent of mixing of the connate water endmember with the low-salinity HU1 groundwaters. These three wells
832 PG81, PG10, and PG76 are found along the important right strike-slip NGFZ_ NW-SE_S limiting the GTP to the north
833 (Figure 1Sa). PG10 and PG76 wells are situated close to a bend of the NGFZ_ NW-SE_S, locally creating an extensional
834 regime, which favours the uprising of the deep connate waters and their intrusion in HU1, as observed in the pull-apart
835 basins along the Sumatran fault (Muraoka et al., 2010). Also the Galatro thermal springs (35-38°C) are found along the
836 NGFZ_ NW-SE_S, but they represent the discharge of a distinct thermal circuit, as indicated by their Na(Ca)-SO₄(Cl)
837 chemistry and low Cl concentration (167-180 mg/kg).

838 Overall, our study revealed that a direct marine intrusion seems to be very localized despite the high demand for water
839 related to the anthropic activities. The growing water requirement is linked to the increase in industrial settlements and to
840 the land use change that happened during the last two decades. In detail, the main change is due to the intensive and
841 ongoing kiwifruit farming that requires a high-water supply (6000–10,000 m³/ha Holzapfel, 2000) compared to the
842 previous citrus and olive crops, with a water consumption of 4000–6000 m³/ha for citrus crops and almost zero for olive
843 (FAO, 2021). The qualitative and quantitative conditions of the aquifer are kept at acceptable levels due to the high
844 recharge inflow that comes mainly from the Aspromonte massif, where the mean precipitation value is 1274 mm/a. An
845 incipient groundwater-seawater mixing, involving up to 7% seawater, was only recognized in the SW area (Sector *E*), in
846 proximity of Scinà.

847 The obtained results confirm the importance of the multidisciplinary approach to identify and characterize sources and
848 driving mechanisms and produce a reliable conceptual model. These results will provide an important support to the
849 decision-making and management phases of the GTP resources.

850

862

863 **References**

- 864 Abu Al Naeem, M.F., Yusoff, I., Ng, T.F., Maity, J.P., Alias, Y., May, R., Alborsh, H., 2019. A study on the impact of anthropogenic
865 and geogenic factors on groundwater salinization and seawater intrusion in Gaza coastal aquifer. Palestine: An integrated
866 multi-techniques approach. *J. Afr. Earth Sci.* 156, 75–93. <https://doi.org/10.1016/j.jafrearsci.2019.05.006>.
- 867 Alcalá, F. J., Custodio, E., 2008. Using the Cl/Br ratio as a tracer to identify the origin of salinity in aquifers in Spain and Portugal.
868 *Journal of Hydrology* 359(1-2), 189-207.
- 869 Alfarrach, N., Walraevens, K., 2018. Groundwater Overexploitation and Seawater Intrusion in Coastal Areas of Arid and Semi-Arid
870 Regions. *Water* 10(2) 143. <https://doi.org/10.3390/w10020143>
- 871 Alshehri, F., Sattam Almadani, S., El-Sorogy A.S., Alwaqdan, E., Alfaifi H.J., Alharbi, T., 2021. Influence of seawater intrusion and
872 heavy metals contamination on groundwater quality, Red Sea coast, Saudi Arabia. *Mar. Pollut. Bull.* 165, 112094.
873 <https://doi.org/10.1016/j.marpolbul.2021.112094>.
- 874 Apollaro, C., Tripodi, V., Vespasiano, G., De Rosa, R., Dotsika, E., Fuoco, I., Critelli, S., Muto, F., 2019. Chemical, isotopic and
875 geotectonic relations of the warm and cold waters of the Galatro and Antonimina thermal areas, southern Calabria, Italy.
876 *Marine and Petroleum Geology* 109, 469–483. <https://doi.org/10.1016/j.marpetgeo.2019.06.020>
- 877 Apollaro, C., Vespasiano, G., Muto, F., De Rosa, R., Barca, D., Marini, L., 2016. Use of mean residence time of water, flowrate, and
878 equilibrium temperature indicated by water geothermometers to rank geothermal resources. Application to the thermal water
879 circuits of Northern Calabria. *Journal of Volcanology and Geothermal Research* 328, 147–158
- 880 Apollaro, C., Accornero, M., Marini, L., Barca, D., De Rosa, R., 2009 The impact of dolomite and plagioclase weathering on the
881 chemistry of shallow groundwaters circulating in a granodiorite-dominated catchment of the Sila Massif (Calabria, Southern
882 Italy). *Appl. Geochem.* 24, 957–979
- 883 Apollaro, C., Marini, L., Critelli, T., De Rosa, R., Bloise, A., Miriello, D., Catalano, M., Armano, V., 2013. Modeling of the impact of
884 dolomite and biotite dissolution on vermiculite composition in a gneissic shallow aquifer of the Sila Massif (Calabria,
885 Italy). *Applied Geochemistry* 35, 297-311.
- 886 Apollaro, C., Vespasiano, G., De Rosa, R. and Marini, L., 2015. Use of mean residence time and flowrate of thermal waters to evaluate
887 the volume of reservoir water contributing to the natural discharge and the related geothermal reservoir volume. Application
888 to Northern Thailand hot springs. *Geothermics* 58, 62-74.
- 889 Apollaro, C., Caracausi, A., Paternoster, M., Randazzo, P., Aiuppa, A., De Rosa, R., Fuoco, I., Mongelli, G., Muto, F., Vanni, E. and
890 Vespasiano, G., 2020. Fluid geochemistry in a low-enthalpy geothermal field along a sector of southern Apennines chain
891 (Italy). *Journal of Geochemical Exploration*, 219, p.106618.
- 892 Apollaro, C., Di Curzio, D., Fuoco, I., Buccianti, A., Dinelli, E., Vespasiano, G., Castrignanò, A., Rusi, S., Barca, D., Figoli, A.,
893 Gabriele, B., 2022. A multivariate non-parametric approach for estimating probability of exceeding the local natural
894 background level of arsenic in the aquifers of Calabria region (Southern Italy). *Science of the Total Environment* 806, 150345.
- 895 Apollaro, C., Buccianti, A., Vespasiano, G., Vardè, M., Fuoco, I., Barca, D., Bloise, A., Miriello, D., Cofone, F., Servidio, A., De Rosa,
896 R., 2019. Comparative geochemical study between the tap waters and the bottled mineral waters in Calabria (Southern Italy)
897 by compositional data analysis (CoDA) developments. *Applied Geochemistry* 107, pp.19-33.
- 898 Appelo C., Postma D., 2005. *Geochemistry, Groundwater and Pollution*, 2nd ed.. Balkema, Rotterdam.
899 <http://dx.doi.org/10.1201/9781439833544>
- 900 [dataset] ARPACAL, Centro Funzionale Multirischi. Consultazione banca dati storici.
901 http://www.cfd.calabria.it/index.php?option=com_wrapper&view=wrapper&Itemid=41
- 902 Awaleh, M. O., Boschetti, T., Adaneh, A. E., Chirdon, M. A., Ahmed, M. M., Dabar, O. A., Soubaneh Y.D. , Egueh N.M., Kawalich
903 A.D., Kadieh I.H., Chaheire, M., 2022. Origin of nitrate and sulfate sources in volcano-sedimentary aquifers of the East
904 Africa Rift System: An example of the Ali-Sabieh groundwater (Republic of Djibouti). *Science of The Total Environment*,
905 804, 150072.
- 906 Barberi, S., Sambito, M., Neduzha, L., Severino, A., 2021. Pollutant Emissions in Ports: A Comprehensive Review. *Infrastructures* 6,
907 114. <https://doi.org/10.3390/infrastructures6080114>

908 Barker, W. W., Welch, S. A., Chu, S., & Banfield, J. F., 1998. Experimental observations of the effects of bacteria on aluminosilicate
909 weathering. *American Mineralogist* 83(11), 1551-1563.

910 Barlow M.P., 2003. Groundwater in Freshwater-Saltwater Environments of the Atlantic Coast. U.S. Geological Survey Circular,
911 Reston, GA, USA..

912 Barlow, P. M., Reichard, E. G., 2010. Saltwater intrusion in coastal regions of North America. *Hydrogeology Journal* 18(1), 247-260.

913 Barrocu, G., 2003. Seawater intrusion in coastal aquifers of Italy. *Tecnología de la Intrusión de Agua de Mar en Acuíferos Costeros:*
914 *Países Mediterráneos*. Instituto Geológico y Minero de España, Madrid.

915 Bellafore, D., Ferrarin, C., Maicu, F., Manfè, G., Lorenzetti, G., Umgiesser, G., Zaggia L., Valle Levinson A., 2021. Saltwater intrusion
916 in a Mediterranean delta under a changing climate. *J. Geophys. Res. Oceans* 126, e2020JC016437.
917 <https://doi.org/10.1029/2020JC016437>

918 Berner, E.K., Berner, R.A., 1996. *Global Environment: Water, Air and Geochemical Cycles*. Prentice Hall, Upper Saddle River.

919 Bocanegra, E.; Da Silva, G.C.; Custodio, E.; Manzano, M.; Montenegro, S. State of knowledge of coastal aquifer management in South
920 America. *Hydrogeol. J.* 2010, 18, 261–267.

921 Bompoti, N., Chrysochoou, M., Dermatas, D., 2015. Geochemical characterization of Greek ophiolitic environments using statistical
922 analysis. *Environ. Process* 2, 5-21. doi:10.1007/s40710-015-0097-z.

923 Boughriba, M., Jilali, A., 2018. Climate change and modeling of an unconfined aquifer: the Triffa plain, Morocco. *Environment,*
924 *Development and Sustainability* 20(5), 2009-2026.

925 Boumaiza, L., Chesnaux, R., Drias, T., Walter, J., Huneau, F., Garel, E., Knoeller, K., Stumpp, C., 2020. Identifying groundwater
926 degradation sources in a Mediterranean coastal area experiencing significant multi-origin stresses. *Science of The Total*
927 *Environment* 746, 141203.

928 Brozzo, G., Accornero, M., Marini, L., (2011). The alluvial aquifer of the Lower Magra Basin (La Spezia, Italy): conceptual
929 hydrogeochemical–hydrogeological model, behavior of solutes, and groundwater dynamics. *Carbonates and Evaporites*
930 26(3), 235-254.

931 Bruni, J., Canepa, M., Cipolli, F., Marini, L., Ottonello, G., Vetuschì Zuccolini, M., Chiodini, G., Cioni, R., Longinelli, A., 2002.
932 Irreversible water-rock mass transfer accompanying the generation of the neutral, Mg-HCO₃ and high-pH, Ca-OH spring
933 waters of the Genoa province, Italy. *Applied Geochemistry* 17, 455-474.

934 Capaccioni, B., Didero, M., Paletta, C., and Didero, L., 2005. Saline intrusion and refreshing in a multilayer coastal aquifer in the
935 Catania Plain (Sicily, Southern Italy): dynamics of degradation processes according to the hydrochemical characteristics of
936 groundwaters. *J. Hydrol.* 307(1-4), 1-16.

937 Cardoso, P., 1993. Saline water intrusion in Mexico, in: Brebbia, C.A. (Ed.), *Water Pollution*, vol. 2, Transactions on Ecology and the
938 Environment. WIT Press, Southampton.

939 Carreira, P. M., Marques, J. M., Nunes, D., 2014. Source of groundwater salinity in coastline aquifers based on environmental isotopes
940 (Portugal): Natural vs. human interference. A review and reinterpretation. *J. Appl. Geochem.* 41, 163–175. <https://doi.org/10.1016/j.apgeochem.2013.12.012>.

942 Carrión-Mero, P., Montalván, F.J., Morante-Carballo, F., Heredia, J., Elorza, F.J., Solórzano, J., Aguilera, H., 2021. Hydrochemical
943 and Isotopic Characterization of the Waters of the Manglaralto River Basin (Ecuador) to Contribute to the Management of
944 the Coastal Aquifer. *Water* 13, 537. <https://doi.org/10.3390/w13040537>

945 Cartwright, I., Weaver, T. R., Fifield, L. K., 2006. Cl/Br ratios and environmental isotopes as indicators of recharge variability and
946 groundwater flow: an example from the southeast Murray Basin, Australia. *Chemical Geology* 231(1-2), 38-56.

947 Chen, Q., Jia, C., Wei, J., Dong, F., Yang, W., Hao, D., Jia Z. & Ji, Y., 2020. Geochemical process of groundwater fluoride evolution
948 along global coastal plains: Evidence from the comparison in seawater intrusion area and soil salinization area. *Chemical*
949 *Geology* 552, 119779.

950 Chien, N. P., & Lautz, L. K., 2018. Discriminant analysis as a decision-making tool for geochemically fingerprinting sources of
951 groundwater salinity. *Science of the Total Environment* 618, 379-387.

952 Cianflone, G., Dominici, R., Viscomi, A., 2015a. Potential recharge estimation of the Sibari Plain aquifers (southern Italy) through a
953 new GIS procedure. *Geographia Technica* 10 (1), 8–18.

954 Cianflone, G., Vespasiano, G., De Rosa, R., Dominici, R., Apollaro, C., Vaselli, O., Pizzino, L., Tolomei, C., Capecchiacci, F., Polemio,
955 M., 2021. Hydrostratigraphic Framework and Physicochemical Status of Groundwater in the Gioia Tauro Coastal Plain
956 (Calabria—Southern Italy). *Water* 13, 3279. <https://doi.org/10.3390/w13223279>.

957 Cianflone, G., Vespasiano, G., Tolomei, C., De Rosa, R., Dominici, R., Apollaro, C., Walraevens, K., Polemio, M., 2022. Different
958 Ground Subsidence Contributions Revealed by Integrated Discussion of Sentinel-1 Datasets, Well Discharge, Stratigraphical
959 and Geomorphological Data: The Case of the Gioia Tauro Coastal Plain (Southern Italy). *Sustainability* , 2926.
960 <https://doi.org/10.3390/su14052926>.

961 Cioni R., Marini L., 2020. A Thermodynamic Approach to Water Geothermometry. Springer Geochemistry Series, Springer Cham,
962 Switzerland. <https://doi.org/10.1007/978-3-030-54318-1>.

963 Cipolli, F., Gambardella, B., Marini, L., Ottonello, G., Vetuschì Zuccolini, M., 2004. Geochemistry of high-pH waters from
964 serpentinites of the Gruppo di Voltri (Genoa, Italy) and reaction path modeling of CO₂ sequestration in serpentinite aquifers.
965 *Applied Geochemistry* 19, 787-802.

966 Cirrincione, R., Fazio, E., Fiannacca, P., Ortolano, G., Pezzino, A., Punturo, R., 2015. The Calabria-Peloritani Orogen, a composite
967 terrane in Central Mediterranean; its overall architecture and geodynamic significance for a pre-Alpine scenario around the
968 Tethyan basin. *Periodico di Mineralogia* 84(3B), 701-749.

969 Corniello, A., Cardellicchio, N., Cavuoto, G., Cuoco, E., Ducci, D., Minissale, A., Mussi, M., Petruccione, E., Pelosi, N., Rizzo, E.,
970 Polemio, M., Tamburino, S., Tedesco, D., Tiano, P., Iorio, M., 2015. Hydrogeological characterization of a geothermal
971 system: The case of the thermo-mineral area of Mondragone (Campania, Italy). *International Journal of Environmental*
972 *Research* 9(2), 523–534. <https://doi.org/10.22059/IJER.2015.926>

973 Craig, H., 1961. Isotopic variations in meteoric waters. *Science*,133(3465), 1702-1703.

974 Custodio, E., 2010. Coastal aquifers of Europe: An overview. *Hydrogeol. J.* 18, 269–280.

975 Davis, S. N., Cecil, L. D., Zreda, M., Moysey, S., 2001. Chlorine-36, bromide, and the origin of spring water. *Chemical Geology*
976 179(1-4), 3-16.

977 De Montety, V., Radakovitch, O., Vallet-Coulomb, C., Blavoux, B., Hermitte, D., Valles, V., 2008. Origin of groundwater salinity
978 and hydrogeochemical processes in a confined coastal aquifer: case of the Rhône delta (Southern France). *Applied*
979 *Geochemistry* 23(8), 2337-2349.

980 De Rosa, R., Dominici, R., Donato, P., Barca D., 2008. Widespread syn-eruptive volcanoclastic deposits in the Pleistocenic basins of
981 South-Western Calabria. *Journal of Volcanology and Geothermal Research* 177, 155-169.

982 Dharmagunawardhane, H.A., Malaviarachchi, K.P.S., Burgess, W., 2016. Fluoride content of minerals in gneissic rocks at an area of
983 endemic dental fluorosis in Sri Lanka: estimates from combined petrographic and electron microprobe analysis. *Ceylon J.*
984 *Sci* 45 (1), 57–66.

985 Drever J. I., 1988 The geochemistry of natural waters. Prentice hall, Englewood Cliffs.

986 Durov, S.A., 1948. Natural Waters and Graphic Representation of Their Composition. *Doklady Akademii Nauk SSSR* 59, 87-90.

987 Edmunds, W. M., 2001. Palaeowaters in European coastal aquifers—the goals and main conclusions of the PALAEAUX project.
988 *Geological Society London Special Publications* 189(1), 1-16.

989 El Halimi, N., Chaouni Alia, A., Beeuwsaert, E., Walraevens, K., 2001. Hydrogeological and Geophysical Investigation for
990 Characterizing the Groundwater Reservoir in Saidia Plain (north-eastern Morocco), in: Walraevens, K. (Ed.), *Development*
991 *of Water Resource Management Tools for Problems of Seawater Intrusion and Contamination of Fresh-Water Resources in*
992 *Coastal Aquifers*. Ghent University, Ghent.

993 Fadili, A., Mehdi, K., Riss, J., Najib, S., Makan, A., Boutayab, K., 2015. Evaluation of groundwater mineralization processes and
994 seawater intrusion extension in the coastal aquifer of Oualidia, Morocco: hydrochemical and geophysical approach. *Arab. J.*
995 *Geosci.* 8, 8567–8582

996 Fehdi, C., Rouabhia, A., Baali, F., Boudoukha, A., 2009. The hydrogeochemical characterization of Morsott-El Aouinet aquifer,
997 Northeastern Algeria. *Environ. Geol.* 58, 1611–1620. doi:10.1007/s00254-008-1667-4.

998 Felisa, G., Ciriello, V., Di Federico, V., 2013. Saltwater Intrusion in Coastal Aquifers: A Primary Case Study along the Adriatic Coast
999 Investigated within a Probabilistic Framework. *Water* 5, 1830-1847; doi:10.3390/w5041830

1000 Fidelibus, M. D. (2003). Environmental tracing in coastal aquifers: old problems and new solutions, in: Lopez-Geta, J.A., de Dios
1001 Gomez, J., de la Orden, J.A., Ramos, G., Rodriguez, L. (Eds.), Coastal aquifers intrusion technology: Mediterranean
1002 countries. IGME, Madrid, pp. 79-111.

1003 Figoli, A., Fuoco, I., Apollaro, C., Chabane, M., Mancuso, R., Gabriele, B., De Rosa, R., Vespasiano, G., Barca, D., Criscuoli, A.,
1004 2020. Arsenic-contaminated groundwaters remediation by nanofiltration. Separation and Purification Technology 238,
1005 116461.

1006 Flury, M., Prapitz, A., 1993. Bromide in the natural environment: occurrence and toxicity. J. Environ. Qual. 22, 747–758.

1007 Food and Agriculture Organization (FAO). Land & Water. [https://www.fao.org/land-water/databases-and-software/crop-](https://www.fao.org/land-water/databases-and-software/crop-information/olive/en/)
1008 [information/olive/en/](https://www.fao.org/land-water/databases-and-software/crop-information/olive/en/) (accessed on 2 October 2021).

1009 Franceschini, F., Signorini, R., 2016. Seawater intrusion via surface water vs. deep shoreline salt-wedge: A case history from the Pisa
1010 coastal plain (Italy). Groundwater for Sustainable Development 2, 73-84.

1011 Fuoco I., Marini L., De Rosa R., Figoli A., Gabriele B., Apollaro C., 2022. Use of reaction path modelling to investigate the evolution
1012 of water chemistry in shallow to deep crystalline aquifers with a special focus on fluoride. Science of the Total Environment
1013 830, 154566

1014 Galli P., Bosi, V., 2002. Paleoseismology along the Cittanova fault: Implications for seismotectonics and earthquake recurrence in
1015 Calabria (southern Italy). J. Geophys. Res. 107, 2044. <https://doi.org/10.1029/2001JB000234>.

1016 Galli, P., Peronace, E., 2015. E. Low slip rates and multimillennial return times for Mw 7 earthquake faults in southern Calabria (Italy).
1017 Geophys. Res. Lett. 42, 5258-5265. <https://doi.org/10.1002/2015GL064062>.

1018 Ganyaglo, SY., Osae, S., Akiti, T., Armah, T., Vitvar, T., Ito, M., Otoo, I.A., 2017. Application of geochemical and stable isotopic
1019 tracers to investigate groundwater salinity in the Ochi-Narkwa Basin Ghana. Hydrol Sci J. 62 (8), 1301-1316.

1020 Garrels, R.M., 1968. Genesis of some ground waters from igneous rocks, in: Abelson, P.H. (Ed.), Researches in Geochemistry vol. 2.
1021 Wiley, New York, pp. 406–420.

1022 Gat, J. R. (1996). Oxygen and hydrogen isotopes in the hydrologic cycle. Annual Review of Earth and Planetary Sciences, 24, 225-
1023 262.

1024 Geoportale Nazionale, Ministero dell’Ambiente. Modello digitale del terreno - 20 metri.
1025 http://wms.pcn.minambiente.it/wcs/dtm_20m (accessed 13 March 2022).

1026 Giambastiani, B.M.S., Kidanemariam, A., Dagnew, A., Antonellini, M., 2021. Evolution of Salinity and Water Table Level of the
1027 Phreatic Coastal Aquifer of the Emilia Romagna Region (Italy). Water, 13, 372. <https://doi.org/10.3390/w13030372>

1028 Giggenbach, W. F., Matsuo, S., 1991. Evaluation of results from Second and Third IAVCEI field workshops on volcanic gases, Mt
1029 Usu, Japan, and White Island, New Zealand. Applied Geochemistry 6(2), 125-141.

1030 Giménez-Forcada, E., Bencini, A., Pranzini, G., 2010. Hydrogeochemical considerations about the origin of groundwater salinization
1031 in some coastal plains of Elba Island (Tuscany, Italy). Environ. Geochem. Health 32, 243–257

1032 Giménez- Forcada, E., & Sánchez San Román, F. J., 2015. An Excel Macro to Plot the HFE- Diagram to Identify Sea Water Intrusion
1033 Phases. Groundwater 53(5), 819-824.

1034 Giustini, F., Brilli, M., Patera, A., 2016. Mapping oxygen stable isotopes of precipitation in Italy. Journal of Hydrology: Regional
1035 Studies 8, 162-181.

1036 Grassi, S., Cortecchi, G., Squarci, P., 2007. Groundwater resource degradation in coastal plains: the example of the Cecina area
1037 (Tuscany–Central Italy). Applied Geochemistry 22(11), 2273-2289.

1038 Han, D., Currell, M. J., Guo, H., 2021. Controls on distributions of sulphate, fluoride, and salinity in aquitard porewater from the North
1039 China Plain: Long-term implications for groundwater quality. Journal of Hydrology 603, 126828.

1040 Helgeson, H. C., 1968. Evaluation of irreversible reactions in geochemical processes involving minerals and aqueous solutions—I.
1041 Thermodynamic relations. Geochimica et Cosmochimica Acta 32(8), 853-877.

1042 Holzapfel, E., Merino, R., Mariño, M., Matta, R., 2000. Water production functions in kiwi. Irrig. Sci. 19, 73–79.

1043 ISPRA, 2016. Foglio 590-Taurianova. In Carta Geologica d'Italia alla Scala 1:50.000. Istituto Superiore per la Protezione e la Ricerca
1044 Ambientale, Roma.

1045 Iyalomhe, F., Rizzi, J., Pasini, S., Torresan, S., Critto, A., Marcomini, A., 2015. Regional Risk Assessment for climate change impacts
1046 on coastal aquifers. *Science of the Total Environment* 537, 100–114. <http://dx.doi.org/10.1016/j.scitotenv.2015.06.111>.

1047 Jacques, E., Monaco, C., Taponnier, P., Tortorici, L., Winter, T., 2001. Faulting and earthquake triggering during the 1783 Calabria
1048 seismic sequence. *Geophys. J. Int.* 147, 499–516.

1049 Kammoun, S., Trabelsi, R., Re, V., Zouari, K., 2021. Coastal aquifer salinization in semi-arid regions: the case of Grombalia (Tunisia).
1050 *Water* 13(2), 129.

1051 Kushnir, J., 1980. The coprecipitation of strontium, magnesium, sodium, potassium and chloride ions with gypsum. An experimental
1052 study. *Geochim. Cosmochim. Ac.* 44(10), 1471–1482.

1053 Kennedy V. C., Zellweger G. W., Jones, B. F., 1974. Filter pore- size effects on the analysis of Al, Fe, Mn, and Ti in water. *Water*
1054 *Resources Research* 10(4), 785–790.

1055 Khairy, H., Janardhana, M., 2013. Hydrogeochemical features of groundwater of semiconfined coastal aquifer in Amol–Ghaemshahr
1056 plain, Mazandaran Province, Northern Iran. *Environ. Monit. Assess.* 185, 9237–9264

1057 Langman, J.B.; Ellis, A.S., 2010. A multi-isotope (δD , $\delta^{18}O$, $87Sr/86Sr$, and $\delta^{11}B$) approach for identifying saltwater intrusion and
1058 resolving groundwater evolution along the Western Caprock Escarpment of the Southern High Plains, New Mexico. *Applied*
1059 *Geochemistry* 25, 159–174.

1060 Laxen, D. P., Chandler, I. M., 1982. Comparison of filtration techniques for size distribution in freshwaters. *Analytical Chemistry*
1061 54(8), 1350–1355.

1062 Li, C., Gao, X., Liu, Y., Wang, Y., 2019. Impact of anthropogenic activities on the enrichment of fluoride and salinity in groundwater
1063 in the Yuncheng Basin constrained by Cl/Br ratio, $\delta^{18}O$, δ^2H , $\delta^{13}C$ and δ^7Li isotopes. *Journal of Hydrology* 579, 124211.

1064 Longhitano, S.G., Chiarella, D., Di Stefano, A., Messina, C., Sabato, L., Tropeano, M., 2012. Tidal signatures in Neogene to Quaternary
1065 mixed deposits of southern Italy straits and bays. *Sediment. Geol.* 279, 74–96. <https://doi.org/10.1016/j.sedgeo.2011.04.019>.

1066 Longinelli, A., Selmo, E., 2003. Isotopic composition of precipitation in Italy: a first overall map. *Journal of Hydrology* 270(1–2), 75–
1067 88.

1068 Loreto, M. F., Düşünür-Doğan, D., Üner, S., İşcan-Alp, Y., Ocakoğlu, N., Cocchi, L., Muccini F., Giordano P., Ligi, M., 2019. Fault-
1069 controlled deep hydrothermal flow in a back-arc tectonic setting, SE Tyrrhenian Sea. *Scientific reports* 9(1), 1–14.

1070 Mallick, J., Singh, C.K., AlMesfer, M.K., Kumar, A., Khan, R.A., Islam, S.; Rahman, A., 2018. Hydro-geochemical assessment of
1071 groundwater quality in Aseer Region, Saudi Arabia. *Water* 10, 1–14, doi:10.3390/w10121847.

1072 Marconi, V., Antonellini, M., Balugani, E., Dinelli, E., 2011. Hydrogeochemical characterization of small coastal wetlands and forests
1073 in the Southern Po plain (Northern Italy). *Ecohydrology* 4, 597–607

1074 Marini, L., Bonaria, V., Guidi, M., Hunziker, J.C., Ottonello, G., Zuccolini, M.V., 2000. Fluid geochemistry of the Acqui Terme-
1075 Visone geothermal area (Piemonte, Italy). *Applied Geochemistry*, 15(7), 917–935.

1076 Marini L., 2006. Geological Sequestration of Carbon Dioxide - Thermodynamics, Kinetics, and Reaction Path Modeling. Elsevier
1077 Science, Amsterdam.

1078 Martínez, M. L., Intralawan, A., Vázquez, G., Pérez-Maqueo, O., Sutton, P., Landgrave, R., 2007. The coasts of our world: Ecological,
1079 economic and social importance. *Ecological economics* 63(2–3), 254–272.

1080 Masciopinto, C., 2006. Simulation of coastal groundwater remediation: The case of Nardò fractured aquifer in Southern Italy. *Environ.*
1081 *Model. Softw.* 21, 85–97.

1082 Mastrocicco, M., Colombani, N., 2020. The Issue of Groundwater Salinization in Coastal Areas of the Mediterranean Region: A
1083 Review. *Water* 13, 90. <https://doi.org/10.3390/w13010090>.

1084 Mastrocicco, M., Gervasio, M. P., Busico, G., Colombani, N., 2021. Natural and anthropogenic factors driving groundwater resources
1085 salinization for agriculture use in the Campania plains (Southern Italy). *Science of the Total Environment* 758, 144033.

1086 Matiatos, I., Paraskevopoulou, V., Lazogiannis, K., Botsou, F., Dassenakis, M., Ghionis, G., Alexopoulos, J. D., Poulos, S. E., 2018.
1087 Surface–ground water interactions and hydrogeochemical evolution in a fluvio-deltaic setting: The case study of the Pinios
1088 River delta. *Journal of Hydrology* 561, 236–249.

1089 McGranahan, G., Balk, D., Anderson, B., 2007. The rising tide: assessing the risks of climate change and human settlements in low
1090 elevation coastal zones. *Environment and urbanization* 19(1), 17–37.

1091 Menció, A., Mas-Pla, J., Otero, N., Regàs, O., Boy-Roura, M., Puig, R., Bach, J., Domènech, C., Zamorano, M., Brusi, D., Folch, A.,
1092 2016. Nitrate pollution of groundwater; all right..., but nothing else?. *Science of the Total Environment*, 539, 241-251.

1093 Mendizabal, I., Baggelaar, P.C., Stuyfzand, P.J., 2012. Hydrochemical trends for public supply well fields in The Netherlands (1898–
1094 2008), natural backgrounds and upscaling to groundwater bodies. *J. Hydrol.* 450 (451), 279–292.

1095 Menichini, M., Doveri, M., 2020. Modelling tools for quantitative evaluations on the Versilia coastal aquifer system (Tuscany, Italy)
1096 in terms of groundwater components and possible effects of climate extreme events. *Acque Sotterranee . Italian Journal of*
1097 *Groundwater* 9(3), 35–44. DOI: 10.7343/as-2020-475

1098 Mjemah, I.C.; Van Camp, M.; Walraevens, K., 2009. Groundwater exploitation and hydraulic parameter estimation for a Quaternary
1099 aquifer in Dar-es-Salaam, Tanzania. *J. Afr. Earth Sci.* 55, 134–146.

1100 Mollema, P. N., Antonellini, M., Dinelli, E., Gabbianelli, G., Greggio, N., Stuyfzand, P. J., 2013. Hydrochemical and physical processes
1101 influencing salinization and freshening in Mediterranean low-lying coastal environments. *Applied Geochemistry* 34, 207-
1102 221.

1103 Monaco, C., Tortorici, L., 2000. Active faulting in the Calabrian arc and eastern Sicily. *J. Geodyn.* , 29, 407-424.

1104 Mtoni, Y., Chikira, ICM., Bakundukize, C., Van Camp, M., Martens, K., Walraevens, K., 2013. Saltwater intrusion and nitrate pollution
1105 in the coastal aquifer of Dar es Salaam Tanzania. *Environ. Earth Sci.* 70, 1091–1111. [https://doi.org/10.1007/s12665-012-](https://doi.org/10.1007/s12665-012-2197-7)
1106 [2197-7](https://doi.org/10.1007/s12665-012-2197-7)

1107 Muraoka, H., Takahashi, M., Sundhoro, H., Dwipa, S., Soeda, Y., Momit, M., Shimada K., 2010. Geothermal Systems Constrained by
1108 the Sumatran Fault and Its Pull-Apart Basins in Sumatra, Western Indonesia, in: *Proceedings World Geothermal Congress*,
1109 Bali, Indonesia, pp. 25-29.

1110 Muzzillo, R., Zuffianò, L.E., Rizzo, E., Canora, F., Capozzoli, L., Giampaolo, V., De Giorgio, G., Sdao, F., Polemio, M., 2021.
1111 Seawater Intrusion Proneness and Geophysical Investigations in the Metaponto Coastal Plain (Basilicata, Italy). *Water* 13(1),
1112 53. <https://doi.org/10.3390/w13010053>

1113 Nasher, G., Al-Sayyaghi, A., Al-Matary, A., 2013. Identification and evaluation of the hydrogeochemical processes of the lower part
1114 of Wadi Siham catchment area, Tihama plain, Yemen. *Arab. J. Geosci.* 6, 2131–2146.

1115 Nisi, B., Vaselli, O., Elio, J., Giannini, L., Tassi, F., Guidi, M., Darrah, T.H., Maletic, E.L., Delgado Huertas, A., Marchionni, S., 2019.
1116 The Campo de Calatrava Volcanic Field (central Spain): Fluid geochemistry in a CO₂-rich area. *Applied Geochemistry* 102,
1117 153-170. doi: <https://doi.org/10.1016/j.apgeochem.2019.01.011>.

1118 Nisi, B., Vaselli, O., Huertas, A. D., Tassi, F., 2013. Dissolved nitrates in the groundwater of the Cecina Plain (Tuscany, Central-
1119 Western Italy): Clues from the isotopic signature of NO₃. *Applied Geochemistry* 34, 38-52.

1120 Nisi, B., Vaselli, O., Taussi, M., Doveri, M., Menichini, M., Cabassi, J., Raco, B., Botteghi, S., Mussi, M., Masetti, G., 2022.
1121 Hydrogeochemical surveys of shallow coastal aquifers: A conceptual model to set-up a monitoring network and increase the
1122 resilience of a strategic groundwater system to climate change and anthropogenic pressure. *Applied Geochemistry* 142,
1123 <https://doi.org/10.1016/j.apgeochem.2022.105350>

1124 Nollet L.M.L., De Gelder L.S.P., 2007. *Handbook of Water Analysis*, second ed. Taylor & Francis,
1125 London. <https://doi.org/10.1201/9781420006315>

1126 Nordstrom, D.K., 1997. Thermochemical redox equilibria of ZoBell's solution. *Geochim. Cosmochim. Acta* 41, 1835–1841.

1127 Palandri, J.L., Reed, M.H., 2001. Reconstruction of in situ composition of sedimentary formation waters. *Geochim. Cosmochim. Acta*
1128 65, 1741–1767.

1129 Parkhurst, D. L., Appelo, C. A. J., 1999. User's guide to PHREEQC (Version 2): A computer program for speciation, batch-reaction,
1130 one-dimensional transport, and inverse geochemical calculations, in: *Water-resources investigations report.. U.S. Geological*
1131 *Survey* 99-4259.

1132 Pastorelli, S., Marini, L., Hunziker, J.C., 2001. Chemistry, isotope values (δD , $\delta^{18}\text{O}$, $\delta^{34}\text{S}_{\text{SO}_4}$) and temperatures of the water inflows in
1133 two Gotthard tunnels, Swiss Alps. *Applied Geochemistry* 16, 633- 649.

1134 Petalas, C., Pisinaras, V., Gemitzi, A., Tsihrintzis, V. A., Ouzounis, K., 2009. Current conditions of saltwater intrusion in the coastal
1135 Rhodope aquifer system, northeastern Greece. *Desalination* 237(1-3), 22-41.

1136 Pfahl, S., Sodemann, H., 2014. What controls deuterium excess in global precipitation?. *Climate of the Past* 10(2), 771-781.

1137 Phocaides A., 2000. Handbook on Pressurized Irrigation Techniques. FAO, Roma..

1138 Pierre, C., 1999. The oxygen and carbon isotope distribution in the Mediterranean water masses. *Marine Geology* 153(1-4), 41-55.

1139 Pizzino, L., Burrato, P., Quattrocchi, F., Valensise, G., 2004. Geochemical signatures of large active faults: The example of the 5
1140 February 1783, Calabrian earthquake (southern Italy). *J. Seismol.* 8, 363–380.

1141 Polemio, M., Walraevens, K., 2019. Recent Research Results on Groundwater Resources and Saltwater Intrusion in a Changing
1142 Environment. *Water* 11 (6), 1118. <https://doi.org/10.3390/w11061118>

1143 Polemio, M., Zuffianò, L. E., 2020. Review of utilization management of groundwater at risk of salinization. *J. Water Resour. Plan.*
1144 *Manag* 146(9), 03120002.

1145 Randazzo, P., Caracausi, A., Aiuppa, A., Cardellini, C., Chiodini, G., Apollaro, C., Paternoster M., Rosiello, A., Vespasiano, G., 2022.
1146 Active degassing of crustal CO₂ in areas of tectonic collision: A case study from the Pollino and Calabria sectors (Southern
1147 Italy). *Front. Earth Sci.* 10, 946707. doi: 10.3389/feart.2022.946707

1148 Reimann, C., De Caritat, P. 1998. Chemical elements in the environment. Springer, New York, Inc., New York, 397 p

1149 Rickard D., 2012. Sulfidic sediments and sedimentary rocks. Elsevier, Amsterdam.

1150 Rickard, D., Musmann, M., Steadman, J. A., 2017. Sedimentary sulfides. *Elements* 13(2), 117-122.

1151 Sarker, M. M. R., Van Camp, M., Hossain, D., Islam, M., Ahmed, N., Karim, M. M., Bhuiyan Md. A. Q., Walraevens, K., 2021.
1152 Groundwater salinization and freshening processes in coastal aquifers from southwest Bangladesh. *Science of The Total*
1153 *Environment* 779, 146339.

1154 Schmidt, G. A., 1999. Forward modeling of carbonate proxy data from planktonic foraminifera using oxygen isotope tracers in a global
1155 ocean model. *Paleoceanography* 14(4), 482-497.

1156 Seal, R. R., 2006. Sulfur isotope geochemistry of sulfide minerals. *Reviews in mineralogy and geochemistry* 61(1), 633-677.

1157 Stenni, B., Nichetto, P., Bregant, D., Scarazzato, P., Longinelli, A., 1995. The Delta-O-18 signal of the northward flow of
1158 mediterranean waters in the adriatic sea. *Oceanologica acta* 18(3), 319-328.

1159 Stewart, F. H., 1963. Marine evaporites, in *Data of geochemistry* (6th ed.): U.S. Geological Survey Professional Paper 440-Y, 52 p.

1160 Steyl, G., Dennis, I., 2010. Review of coastal-area aquifers in Africa. *Hydrogeol. J.* 18, 217–225.

1161 Stumm W., Morgan J.J., 1996. Aquatic chemistry: chemical equilibria and rates in natural waters, third ed. Wiley, New York

1162 Stuyfzand, P. J., 1989. A new hydrochemical classification of water types. *Iahs Publ.* 182, 89-98.

1163 Stuyfzand, P., 1986. A New Hydrochemical Classification of water types with examples of application to the Netherlands. *H2O* 19. pp.
1164 562–568.

1165 Stuyfzand, P. J., 2008. Base exchange indices as indicators of salinization or freshening of (coastal) aquifers, in: 20th Salt Water
1166 Intrusion Meeting, Naples, Florida, USA. IFAS Research, Gainesville.

1167 Sudaryanto, Nailly, W., 2018. Ratio of major ions in groundwater to determine saltwater intrusion in coastal areas. *Conf. Ser.: Environ.*
1168 *Earth Sci.* 118, 012021.

1169 Tassi, F., Capeccchiacci, F., Giannini, L., Vougioukalakis, G. Vaselli, O., 2013. Volatile organic compounds (VOCs) in air from Nisyros
1170 Island (Dodecanese Archipelago, Greece): Natural versus anthropogenic sources. *Environ. Poll.* 180,
1171 <https://doi.org/10.1016/j.envpol.2013.05.023>

1172 Taussi, M., Gozzi, C., Vaselli, O., Cabassi, J., Menichini, M., Doveri, M., Romei, M., Ferretti, A., Gambioli, A., Nisi B., 2022.
1173 Contamination Assessment and Temporal Evolution of Nitrates in the Shallow Aquifer of the Metauro River Plain (Adriatic
1174 Sea, Italy) after Remediation Actions. *Int. J. Env. Res. Pub. He.* 19 (19), 12231. <https://doi.org/10.3390/ijerph191912231>

1175 Torres-Martínez, J. A., Mora, A., Mählknecht, J., Daesslé, L. W., Cervantes-Avilés, P. A., Ledesma-Ruiz, R., 2021. Estimation of
1176 nitrate pollution sources and transformations in groundwater of an intensive livestock-agricultural area (Comarca Lagunera),
1177 combining major ions, stable isotopes and MixSIAR model. *Environmental Pollution* 269, 115445.

1178 Tripodi, V., Muto, F., Brutto, F., Perri, F., Critelli, S., 2018. NeogenQuaternary evolution of the forearc and backarc regions between
1179 the Serre and Aspromonte Massifs, Calabria (southern Italy). *Marine and Petroleum Geology* 95, 328-343.

1180 Turekian, K. K., Wedepohl, K. H., 1961. Distribution of the elements in some major units of the earth's crust. *Geological Society of*
1181 *America Bulletin* 72(2), 175-192.

1182 Van Camp, M., Mjemah, I.C., Alfarrak, N., Walraevens, K., 2013. Modeling approaches and strategies for data-scarce aquifers:
1183 Example of the Dar es Salaam aquifer in Tanzania. *Hydrogeol. J.* 21, 341–356.

1184 Van Camp, M., Mtoni, Y.E., Mjemah, I.C., Bakundukize, C., Walraevens, K., 2014. Investigating seawater intrusion due to
1185 groundwater pumping with schematic model simulations: The example of the Dar Es Salaam coastal aquifer in Tanzania. *J.*
1186 *Afr. Earth Sci.* 96, 71–78.

1187 Vandenbohede, A., Lebbe, L., 2012. Groundwater chemistry patterns in the phreatic aquifer of the central Belgian coastal plain. *Applied*
1188 *Geochemistry* 27, 22–36

1189 Vardè, M., Servidio, A., Vespasiano, G., Pasti, L., Cavazzini, A., Di Traglia, M., Rosselli, A., Cofone, F., Apollaro, C., Cairns, W.R.,
1190 Scalabrin, E., 2019. Ultra-trace determination of total mercury in Italian bottled waters. *Chemosphere* 219, 896–913.

1191 Vaselli, O., Lazzaroni, M., Nisi, B., Cabassi, J., Tassi, F., Rappuoli, D., Meloni, F., 2021. Discontinuous Geochemical Monitoring of
1192 the Galleria Italia Circumneutral Waters (Former Hg-Mining Area of Abbadia San Salvatore, Tuscany, Central Italy) Feeding
1193 the Fosso Della Chiusa Creek. *Environment* 8, 15. doi.org/10.3390/environments8020015

1194 Vespasiano, G., Apollaro, C., Marini, L., Dominici, R., Cianflone, G., Romanazzi, A., Polemio, M., De Rosa, R., 2015a.
1195 Hydrogeological and isotopic study of the multi-aquifer system of the Sibari Plain (Calabria, Southern Italy). *Rend. Online*
1196 *Soc. Geol. It.* 39, 134–137

1197 Vespasiano, G., Apollaro, C., De Rosa, R., Muto, F., Larosa, S., Fiebig, J., Mulch, A., Marini, L. (2015b): The Small Spring Method
1198 (SSM) for the definition of stable isotope - elevation relationships in Northern Calabria (Southern Italy). *Appl. Geochem.* 63,
1199 333–346.

1200 Vespasiano, G., Apollaro, C., Muto, F., Dotsika, E., De Rosa, R., Marini, L., 2014. Chemical and isotopic characteristics of the warm
1201 and cold waters of the Luigiane Spa near Guardia Piemontese (Calabria, Italy) in a complex faulted geological framework.
1202 *Applied Geochemistry* 41, 73–88.

1203 Vespasiano, G., Cianflone, G., Cannata, C.B., Apollaro, C., Dominici, R., De Rosa, R., 2016. Analysis of groundwater pollution in the
1204 Sant'Eufemia Plain (Calabria – South Italy). *Italian Journal of Engineering Geology and Environment* 2, 5–15. DOI:
1205 10.4408/IJEGE.2016-02.O-01

1206 Vespasiano, G., Cianflone, G., Romanazzi, A., Apollaro, C., Dominici, R., Polemio, M., De Rosa, R., 2019. A multidisciplinary
1207 approach for sustainable management of a complex coastal plain: The case of Sibari Plain (Southern Italy). *Marine and*
1208 *Petroleum Geology* 109, 740–759. <https://doi.org/10.1016/j.marpetgeo.2019.06.031>.

1209 Vespasiano, G., Marini, L., Muto, F., Auqué, L. F., Cipriani, M., De Rosa, R., Critelli, S., Gimeno, M. J., Blasco, M., Dotsika, E.,
1210 Apollaro, C., 2021a Chemical, isotopic and geotectonic relations of the warm and cold waters of the Cotronei (Ponte
1211 Coniglio), Bruciarello and Repole thermal areas, (Calabria - Southern Italy). *Geothermics* 96,
1212 <https://doi.org/10.1016/j.geothermics.2021.102228>

1213 Vespasiano, G., Muto, F., Apollaro, C., 2021b. Geochemical, Geological and Groundwater Quality Characterization of a Complex
1214 Geological Framework: The Case Study of the Coreca Area (Calabria, South Italy). *Geosciences* 11(3), 121;
1215 <https://doi.org/10.3390/geosciences11030121>

1216 Walraevens, K., Mjemah, I.C., Mtoni, Y., Van Camp, M., 2015. Sources of salinity and urban pollution in the Quaternary sand aquifers
1217 of Dar es Salaam, Tanzania. *J. Afr. Earth Sci.* 102, 149–165.

1218 Werner, A.D. 2010. A review of seawater intrusion and its management in Australia. *Hydrogeol. J.* 18, 281–285.

1219 White, A. F., Bullen, T. D., Vivit, D. V., Schulz, M. S., Clow D. W., 1999. The role of disseminated calcite in the chemical weathering
1220 of granitoid rocks. *Geochim. Cosmochim. Acta* 63 (13–14), 1939–1953

1221 White, A.F., Schulz, M.S., Lowenstern, J.B., Vivit, D.V., Bullen, T.D., 2005. The ubiquitous nature of accessory calcite in granitoid
1222 rocks: Implications for weathering, solute evolution, and petrogenesis. *Geochim. Cosmochim. Acta* 69, 1455–1471.

1223 Wua, X., Zhangb, L., Hub, B.X., Yang Wangb, Y., Xue, Z., 2020. Isotopic and hydrochemical evidence for the salinity origin in the
1224 coastal aquifers of the Pearl River Delta, Guangzhou, China. *J. Contam. Hydrol.* 235, 103732.
1225 doi.org/10.1016/j.jconhyd.2020.103732

1226 Zarei, M., Raeisi, E., Merkel, B.J., 2012. Identifying sources of salinization using hydrochemical and isotopic techniques Konarsiah,
1227 Iran. *Environ. Earth Sci.* 70 (2), 1. DOI: 10.1007/s12665-012-2143-8